

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/BE/19 Row Spacing and Methods of Applying Phorate - Spring Beans

Rothamsted Research

Rothamsted Research (1983) *82/R/BE/19 Row Spacing and Methods of Applying Phorate - Spring Beans* ; Yields Of The Field Experiments 1982, pp 304 - 305 - DOI:

<https://doi.org/10.23637/ERADOC-1-33>

82/R/BE/19

SPRING BEANS

ROW SPACING AND METHODS OF APPLYING PHORATE

Object: To study the effects of rates and methods of applying phorate on the incidence of Sitona and on the yield of s. beans sown on wide or narrow rows - Claycroft.

Sponsors: R. Bardner, D.C. Griffiths, K.E. Fletcher.

Design: 4 randomised blocks of 9 plots.

Whole plot dimensions: 5.33 x 13.7.

Treatments: All combinations of:-

1. PHO RATE Rates of phorate:

PHORATE1	1.7 kg
PHORATE2	2.2 kg

2. PHO METH Methods of applying phorate:

BCAST N	Broadcast and rotary harrowed in to the seedbed, sown in rows 18 cm (7 inches) apart
CDRILL N	Combine drilled with seed, sown in rows 18 cm apart
CDRILL W	Combine drilled with seed, sown in rows 53 cm (21 inches) apart

plus three extra treatments:

EXTRA

O O N	No insecticide, seed sown in rows 18 cm apart
O O W	No insecticide, seed sown in rows 53 cm apart
TE1 BC N	Terbufos at 1.7 kg broadcast and rotary harrowed in to the seedbed, sown in rows 18 cm apart.

Basal applications: Manures: Chalk at 5.0 t. Glyphosate at 1.4 kg in 250 l. Trietazine at 1.0 kg with simazine at 0.14 kg.

Seed: Minden, sown at 270 kg.

Cultivations, etc.: - Glyphosate applied: 4 Nov, 1981. Chalk applied: 10 Nov. Ploughed: 27 Nov. Spring-tine cultivated: 24 Mar, 1982. Terbufos treatment applied: 25 Mar. Remaining broadcast treatments applied, rotary harrowed: 26 Mar. Seed sown: 27 Mar. Trietazine and simazine applied: 1 Apr. Combine harvested: 2 Sept. Previous crops: W. wheat 1980 and 1981.

NOTE: Adult Sitona damage was assessed in late April and mid-May. Larvae were assessed in late June.

82/R/BE/19

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PHO METH	BCAST N	CDRILL N	CDRILL W	MEAN
PHO RATE				
PHORATE1	3.51	3.69	3.76	3.65
PHORATE2	3.74	3.82	3.75	3.77
MEAN	3.62	3.75	3.75	3.71
EXTRA	0 0 N	0 0 W	TE1 BC N	MEAN
	3.73	3.44	3.59	3.59

GRAND MEAN 3.67

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	PHO RATE	PHO METH	PHO RATE PHO METH & EXTRA

SED	0.212	0.123	0.150	0.212

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	24	0.300	8.2

GRAIN MEAN DM% 76.5

PLOT AREA HARVESTED 0.00293