

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/BE/20 Pirimiphos-methyl and Stem Nematode - Spring Beans

Rothamsted Research

Rothamsted Research (1983) *82/R/BE/20 Pirimiphos-methyl and Stem Nematode - Spring Beans* ; Yields Of The Field Experiments 1982, pp 306 - 307 - DOI:

<https://doi.org/10.23637/ERADOC-1-33>

82/R/BE/20

SPRING BEANS

PIRIMIPHOS-METHYL AND STEM NEMATODE

Object: To study the effects of three rates of pirimiphos-methyl applied as a seed dressing on the occurrence of stem nematode, *Sitona* weevils and on the yield of s. beans - Highfield 0 and E VI N.

Sponsors: A.G. Whitehead, J. McEwen, D.P. Yeoman.

Design: 4 randomised blocks of 4 plots.

Whole plot dimensions: 2.03 x 2.13.

Treatments:

PIRIM SD	Pirimiphos-methyl seed dressing (g per tonne of seed):
0	
2	
4	
20	

NOTE: Seed was sown by hand in rows 51 cm apart, seed spaced 5 cm apart in the row.

Basal applications: None.

Seed: Minden.

Cultivations, etc.: - Ploughed: 17 Nov, 1981. Spring-tine cultivated, power harrowed, sown: 26 Mar, 1982. Hand hoed: 12 May, 1 June. Harvested by hand: 7 Sept. Previous crops: S. barley 1980, fallow 1981.

NOTE: Plant counts were made after establishment. Stem nematode symptoms were assessed during the season. Weevil damage was assessed in May.

82/R/BE/20

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PIRIM SD	0	2	4	20	MEAN
	5.83	6.06	4.98	5.02	5.47

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PIRIM SD
-----	-----
SED	0.560

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	9	0.792	14.5

GRAIN MEAN DM% 83.3

PLOT AREA HARVESTED 0.00015