

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/RN/17 Rates of P and K to the Subsoil - Potatoes, S. Barley, S. Beans, W. Wheat

Rothamsted Research

Rothamsted Research (1983) *82/R/RN/17 Rates of P and K to the Subsoil - Potatoes, S. Barley, S. Beans, W. Wheat* ; Yields Of The Field Experiments 1982, pp 83 - 90 - DOI:

<https://doi.org/10.23637/ERADOC-1-33>

82/R/RN/17

RATES OF P AND K TO THE SUBSOIL

Object: To study the effects of a range of rates and frequencies of application of P and K to the subsoil, singly and together, on the yields and nutrient uptakes of a rotation of crops - Meadow.

Sponsors: J. McEwen, A.E. Johnston.

The second year, potatoes, s. barley, s. beans, w. wheat.

For previous year see 81/R/RN/17.

Design: 4 series (for crops) each of 40 plots.

Whole plot dimensions: 3.0 x 14.0.

Treatments to each series:

TREATMNT Extra P and K and primary cultivation tool in autumn 1980 only except on R plots, treatments repeated in autumn 1981:

	P ₂ O ₅ (kg)	K ₂ O(kg)	Tool	
- - -	0	0	Plough	(duplicated)
P6 K6 T	1000	500 to topsoil	"	(")
- - S	0	0 " "	Wye double-digger	(four plots)
- - SR	0	0 " "	"	(duplicated)
P2 - SR	63	0 to subsoil	"	"
P3 - S	125	0 " "	"	"
P4 - S	250	0 " "	"	"
P5 - S	500	0 " "	"	" (duplicated)
P6 - S	1000	0 " "	"	"
- K2 SR	0	31 " "	"	"
- K3 S	0	63 " "	"	"
- K4 S	0	125 " "	"	"
- K5 S	0	250 " "	"	" (duplicated)
- K6 S	0	350 " "	"	"
P1 K1 SR	31	16 " "	"	"
P1 K3 SR	31	63 " "	"	"
P2 K2 SR	63	31 " "	"	"
P3 K1 SR	125	16 " "	"	"
P3 K3 SR	125	63 " "	"	"
P3 K4 S	125	125 " "	"	"
P4 K3 S	250	63 " "	"	"
P4 K4 S	250	125 " "	"	"
P4 K5 S	250	250 " "	"	" (duplicated)
P4 K6 S	250	350 " "	"	"
P5 K4 S	500	125 " "	"	" (duplicated)
P5 K5 S	500	250 " "	"	"
P5 K6 S	500	350 " "	"	"
P6 K4 S	1000	125 " "	"	"
P6 K5 S	1000	250 " "	"	"
P6 K6 S	1000	350 " "	"	"

82/R/RN/17

- NOTES: (1) Subsoiling was done with the Wye double-digger which turns a furrow with a conventional plough share, to a depth of 23 cm, and at the same time rotary cultivates the bottom of the adjacent furrow to a further depth of 15 cm. When applying P and K this was distributed ahead of the rotary cultivator.
- (2) The topsoil PK dressing was equally divided before and after ploughing.
- (3) All plots other than R were conventionally ploughed in autumn 1981.
- (4) The rate of 350 kg K₂O applied was in error for 500 kg K₂O.

Standard applications:

- Potatoes: Manures: (10:10:15 + 4.5 Mg) at 1960 kg. Weedkiller: Metribuzin at 0.98 kg in 900 l. Fungicides: Mancozeb at 1.4 kg in 250 l on four occasions, the first two with the insecticide. Ofurace at 0.12 kg and maneb at 1.2 kg in 250 l on two occasions, the first with the insecticide. Insecticide: Pirimicarb at 0.14 kg.
- S. barley: Manures: (20:10:10) at 450 kg, combine drilled. Weedkillers: Glyphosate at 1.4 kg in 250 l. Dicamba, mecoprop and MCPA (as 'Poly-Farmon' at 5.0 l) in 250 l applied with the fungicide. Fungicide: Tridemorph at 0.53 kg.
- S beans: Weedkillers: Paraquat at 0.6 kg ion in 250 l. Trietazine at 1.0 kg and simazine at 0.14 kg in 250 l. Insecticide: Phorate at 5.6 kg.
- W. wheat: Manures: (0:14:28) at 450 kg. 'Nitro-Chalk' at 480 kg. Weedkillers: Glyphosate at 1.4 kg in 250 l. Mecoprop (as 'Methoxone M' at 5.0 l) and isoproturon at 2.0 kg in 250 l. Fungicide: Propiconazole at 0.12 kg in 250 l on the first occasion at 0.25 kg in 250 l applied with the insecticide on the second occasion. Insecticide: Pirimicarb at 0.14 kg.

Seed: Potatoes: Pentland Crown.

- S. barley: Triumph, seed dressed with ethirimol, sown at 160 kg.
- S. beans: Minden, sown at 270 kg.
- W. wheat: Avalon, sown at 200 kg.

Cultivations, etc.:-

- All crops: Treatments applied by double digger: 30 Oct - 2 Nov, 1981. Ploughed: 3 - 4 Nov.
- Potatoes: Heavy spring-tine cultivated twice: 6 Nov, 1981 and 24 Mar, 1982. NPK Mg applied: 19 Apr. Spiked rotary cultivated, potatoes planted: 20 Apr. Rotary ridged: 10 May. Weedkiller applied: 8 June. Mancozeb with insecticide applied: 14 June, 30 June. Mancozeb applied: 12 July. Ofurace and maneb with insecticide applied: 26 July. Ofurace and maneb applied: 9 Aug. Mancozeb applied: 25 Aug. Haulm mechanically destroyed: 17 Sept. Lifted: 26 Oct.
- S. barley: Heavy spring-tine cultivated twice: 5 Nov, 1981 and 24 Mar, 1982. Rotary harrowed, seed sown: 31 Mar. Dicamba, mecoprop MCPA with tridemorph applied: 10 May. Glyphosate applied: 10 Aug. Combine harvested: 19 Aug.
- S. beans: Heavy spring-tine cultivated: 6 Nov, 1981. Paraquat applied: 23 Mar, 1982. Heavy spring-tine cultivated: 24 Mar. Phorate applied, rotary harrowed, seed sown: 27 Mar. Trietazine and simazine applied: 2 Apr. Combine harvested: 2 Sept.

82/R/RN/17

W. wheat: Heavy spring-tine cultivated twice, PK applied: 5 Nov, 1981. Rotary harrowed, seed sown: 6 Nov. Mecoprop and isoproturon applied: 14 Apr, 1982. N applied: 16 Apr. Propiconazole applied: 26 May. Propiconazole with the insecticide applied: 15 June. Glyphosate applied: 10 Aug. Combine harvested: 21 Aug.

82/R/RN/17 SERIES IV POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	53.7
P6 K6 T	52.9
- - S	56.1
- - SR	54.5
P2 - SR	48.7
P3 - S	50.2
P4 - S	58.3
P5 - S	55.0
P6 - S	50.4
- K2 SR	64.1
- K3 S	52.1
- K4 S	55.4
- K5 S	54.2
- K6 S	56.6
P1 K1 SR	59.3
P1 K3 SR	48.5
P2 K2 SR	59.7
P3 K1 SR	61.3
P3 K3 SR	47.8
P3 K4 S	47.0
P4 K3 S	51.4
P4 K4 S	46.3
P4 K5 S	58.5
P4 K6 S	53.8
P5 K4 S	50.6
P5 K5 S	48.9
P5 K6 S	51.4
P6 K4 S	50.5
P6 K5 S	50.4
P6 K6 S	52.6
MEAN	53.7

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	1.81 MIN REP
	1.56 MAX-MIN

* SED APPLIES ONLY TO - - -, P6 K6 T, - - S, - - SR, P5 - S, - K5 S, P4 K5 S AND P5 K4 S

TREATMNT
 MAX-MIN - - S V ANY OF REMAINDER
 MIN REP ANY OF THE REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	1.81	3.4

82/R/RN/17

SERIES IV POTATOES

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

TREATMNT	
- - -	98.1
P6 K6 T	97.5
- - S	97.3
- - SR	96.8
P2 - SR	96.6
P3 - S	96.8
P4 - S	97.7
P5 - S	96.8
P6 - S	96.8
- K2 SR	98.1
- K3 S	96.3
- K4 S	97.7
- K5 S	97.6
- K6 S	97.3
P1 K1 SR	97.6
P1 K3 SR	97.1
P2 K2 SR	96.1
P3 K1 SR	97.9
P3 K3 SR	96.4
P3 K4 S	96.5
P4 K3 S	97.1
P4 K4 S	96.6
P4 K5 S	96.6
P4 K6 S	97.3
P5 K4 S	97.3
P5 K5 S	96.7
P5 K6 S	96.3
P6 K4 S	96.2
P6 K5 S	97.3
P6 K6 S	94.8
MEAN	97.0

PLOT AREA HARVESTED 0.00210

82/R/RN/17

SERIES I SPRING BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	7.47
P6 K6 T	7.36
- - S	7.37
- - SR	7.32
P2 - SR	7.82
P3 - S	7.52
P4 - S	7.49
P5 - S	7.53
P6 - S	7.61
- K2 SR	7.66
- K3 S	7.45
- K4 S	7.65
- K5 S	7.46
- K6 S	7.40
P1 K1 SR	7.34
P1 K3 SR	7.71
P2 K2 SR	7.71
P3 K1 SR	7.52
P3 K3 SR	7.70
P3 K4 S	7.68
P4 K3 S	7.58
P4 K4 S	7.69
P4 K5 S	7.62
P4 K6 S	7.23
P5 K4 S	7.72
P5 K5 S	7.49
P5 K6 S	7.50
P6 K4 S	7.62
P6 K5 S	7.90
P6 K6 S	7.69
MEAN	7.54

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.195 MIN REP
	0.169 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.195	2.6
GRAIN MEAN DM%	82.7		
PLOT AREA HARVESTED	0.00286		

82/R/RN/17

SERIES II SPRING BEANS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	5.07
P6 K6 T	5.23
- - S	5.48
- - SR	5.22
P2 - SR	5.39
P3 - S	5.09
P4 - S	4.74
P5 - S	4.94
P6 - S	5.27
- K2 SR	5.10
- K3 S	4.79
- K4 S	5.02
- K5 S	5.04
- K6 S	5.36
P1 K1 SR	4.73
P1 K3 SR	5.47
P2 K2 SR	5.33
P3 K1 SR	4.86
P3 K3 SR	5.17
P3 K4 S	5.33
P4 K3 S	4.63
P4 K4 S	5.33
P4 K5 S	4.94
P4 K6 S	4.99
P5 K4 S	5.11
P5 K5 S	5.19
P5 K6 S	5.12
P6 K4 S	5.47
P6 K5 S	5.82
P6 K6 S	5.42
MEAN	5.17

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABL	TREATMNT*
-----	-----
SED	0.432 MIN REP
	0.374 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.432	8.4

GRAIN MEAN DM% 74.8

PLOT AREA HARVESTED 0.00386

82/R/RN/17

SERIES III WINTER WHEAT

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	8.79
P6 K6 T	8.66
- - S	8.64
- - SR	8.75
P2 - SR	8.83
P3 - S	9.03
P4 - S	8.97
P5 - S	8.94
P6 - S	9.28
- K2 SR	9.02
- K3 S	9.09
- K4 S	8.89
- K5 S	8.86
- K6 S	8.70
P1 K1 SR	8.74
P1 K3 SR	8.82
P2 K2 SR	8.97
P3 K1 SR	8.90
P3 K3 SR	8.76
P3 K4 S	8.74
P4 K3 S	8.77
P4 K4 S	8.90
P4 K5 S	8.96
P4 K6 S	8.96
P5 K4 S	8.93
P5 K5 S	9.23
P5 K6 S	9.10
P6 K4 S	8.84
P6 K5 S	8.96
P6 K6 S	8.81
MEAN	8.87

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.226 MIN REP
	0.196 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.226	2.6

GRAIN MEAN DM% 85.2

PLOT AREA HARVESTED 0.00286