

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1982

[Full Table of Content](#)



82/R/CS/10 and 82/W/CS/10 Long Term Liming - S. Oats

Rothamsted Research

Rothamsted Research (1983) *82/R/CS/10 and 82/W/CS/10 Long Term Liming - S. Oats* ; Yields Of The Field Experiments 1982, pp 91 - 93 - DOI: <https://doi.org/10.23637/ERADOC-1-33>

82/R/CS/10 and 82/W/CS/10

LONG TERM LIMING

Object: To study the effects of different amounts of lime on the yields of a sequence of crops. The effects of P are also studied - Rothamsted (R) Sawyers I and Woburn (W) Stackyard C.

Sponsors: S. McGrath, D.P. Stribley.

The 21st year, s. oats.

For previous years see 'Details' 1967, 1973 and 74-81/R&W/CS/10.

Design: 2 randomised blocks of 16 plots.

Whole dimensions: 6.40 x 18.3.

Treatments: All combinations of:-

1. CHALK Ground chalk (tonnes CaCO₃) (total applied 1962-82):

R	W	Rothamsted		Woburn	
		Total 1962-78	1982	Total 1962-78	1982
0	0	0	0	0	0
9	8	7	2	6	2
20	19	15	5	14	5
40	33	30	10	23	10

2. P P fertilizer applied:

	Until 1978	1981	1982
0	None	None	None
P1	K	25 kg P	50 kg P
P2	P	25 kg P	None
P3	P & K	75 kg P	50 kg P

- NOTES: (1) Until 1978 test P & K were applied cumulatively, rates varied with crop. None was applied in 1979 & 1980 (fallow).
 (2) A basal dressing of 120 kg K has been applied since 1981.
 (3) A sub plot test of Mg applied in earlier years has been ignored, but a basal dressing of 100 kg Mg was applied in 1981 and 40 kg Mg in 1982.

Basal applications:

Sawyers I (R): Manures: N at 30 kg as 'Nitro-Chalk' combine drilled, K at 120 kg as muriate of potash, Mg at 40 kg as kieserite.
 Weedkillers: Dicamba, mecoprop and MCPA (as 'Banlene Plus' at 5.0 l) in 250 l.
 Stackyard C (W): Manures: N at 30 kg as 'Nitro-Chalk', K at 120 kg as muriate of potash, Mg at 40 kg as kieserite. Weedkillers: Dicamba, mecoprop and MCPA (as 'Herrisol' at 4.9 l) in 280 l.

Seed: Peniarth, sown at 190 kg (R), 200 kg (W).

Cultivations, etc.:-

Sawyers I (R): Ground chalk, applied: 3-7 Dec, 1981. Ploughed: 3 Feb, 1982. Spring-tine cultivated: 5 Apr. P, K and Mg applied, seed sown: 14 Apr. Weedkillers applied: 26 May. Combine harvested: 26-27 Aug.

82/R/CS/10 and 82/W/CS/10

Stackyard C (W): Ground chalk applied: 25 Nov, 1981. Ploughed: 26 Feb, 1982. P, and K applied: 24 Mar. Mg and N applied, heavy spring-tine cultivated: 25 Mar. Spring-tine cultivated with crumbler attached, seed sown: 29 Mar. Weedkillers applied: 26 May. Combine harvested: 20 Aug.

82/R/CS/10 SAWYERS I (R)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

P	0	P1	P2	P3	MEAN
CHALK					
0	1.21	1.20	1.52	1.74	1.42
9	1.15	1.92	1.47	1.38	1.48
20	1.18	1.22	1.37	1.47	1.31
40	1.13	1.43	1.32	1.65	1.38
MEAN	1.17	1.44	1.42	1.56	1.40

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHALK	P	CHALK P
-----	-----	-----	-----
SED	0.090	0.090	0.180

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	0.180	12.9

GRAIN MEAN DM% 65.8

PLOT AREA HARVESTED 0.00520

82/W/CS/10 STACKYARD C (W)

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

P	0	P1	P2	P3	MEAN
CHALK					
0	1.32	1.79	1.59	1.86	1.64
8	1.58	1.99	1.75	2.08	1.85
19	1.62	1.68	1.86	2.17	1.83
33	1.64	1.77	2.00	1.94	1.84
MEAN	1.54	1.81	1.80	2.01	1.79

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHALK	P	CHALK P

SED	0.096	0.096	0.191

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	15	0.191	10.7

GRAIN MEAN DM% 80.1

PLOT AREA HARVESTED 0.00503