

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/CS/261 Benomyl and Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (1984) *83/R/CS/261 Benomyl and Take-all - W. Wheat* ; Yields Of The Field Experiments 1983, pp 151 - 152 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/CS/261

BENOMYL AND TAKE-ALL

Object: To study the effects of benomyl, applied to the soil with and without a surfactant, and of nuarimol on take-all (*Gaeumannomyces graminis*) and on the yield of w. wheat - New Zealand.

Sponsor: G.L. Bateman.

The third year, w. wheat.

For previous years see 81-82/R/CS/261.

Design: 4 randomised blocks of 7 plots.

Whole plot dimensions: 2.13 x 12.2.

Treatments:

FUNGICIDE	Residues of fungicides and surfactant applied for wheat in 1981 and fresh applications for wheat in 1983:
NONE	None
(BEN A)	Benomyl at 20 kg in autumn 1980
(BEN S)	Benomyl at 20 kg in spring 1981
(NUA A)	Nuarimol at 2.2 kg in autumn 1980
BEN A	Benomyl at 20 kg in autumn 1982
BEN SF A	Benomyl at 20 kg plus a surfactant (an alcohol ethoxylate at 44 l) in autumn 1982
SF A	Surfactant alone in autumn 1982

NOTES: (1) Treatments applied in autumn 1982 were randomly allocated to plots given benomyl, nuarimol or surfactant in spring 1981.
(2) All fresh treatments were applied in 8,800 l on 31 Oct, 1982.

Basal applications: Manures: (0:18:36) at 250 kg. 'Nitro-Chalk' at 630 kg.
Weedkillers: Isoproturon at 2.0 kg with mecoprop at 3.2 kg in 250 l.
Fungicide: Propiconazole at 0.12 kg in 250 l. Growth regulator: Chlormequat at 1.1 kg in 250 l.

Seed: Avalon, seed not treated, sown at 180 kg.

Cultivations, etc.: PK applied: 10 Sept, 1982. Ploughed: 21 Sept.
Spring-tine cultivated, rotary harrowed, seed sown: 15 Oct. N applied: 15 Apr, 1983. Weedkillers applied: 16 Apr. Growth regulator applied: 29 Apr. Fungicide applied: 17 June. Combine harvested: 10 Aug.

NOTE: Take-all and footrots were assessed in May, June and July.

83/R/CS/261

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE	
NONE	9.56
(BEN A)	9.06
(BEN S)	9.84
(NUA A)	10.04
BEN A	9.40
BEN SF A	9.37
SF A	8.96
MEAN	9.46

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGCIDE
-----	-----
SED	0.535

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	18	0.757	8.0

GRAIN MEAN DM% 87.1

PLOT AREA HARVESTED 0.00168