

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1983

[Full Table of Content](#)



## 83/R/CS/271 Aphid Control by Natural Enemies - W. Wheat, Ryegrass

### Rothamsted Research

Rothamsted Research (1984) *83/R/CS/271 Aphid Control by Natural Enemies - W. Wheat, Ryegrass ; Yields Of The Field Experiments 1983*, pp 153 - 154 - DOI:

<https://doi.org/10.23637/ERADOC-1-44>

83/R/CS/271

APHID CONTROL BY NATURAL ENEMIES

Object: To study the effects of early introduction of aphids to ryegrass on subsequent aphid populations on w. wheat - Black Horse I.

Sponsors: W. Powell, A.M. Dewar, N. Wilding, K.E. Fletcher, R. Bardner, C.A. Edwards, J.R. Lofty.

The second year, w. wheat, ryegrass.

For previous year see 82/R/CS/271.

Design: 2 randomised blocks of 4 plots.

Whole plot dimensions: 48 x 48.

Treatments: All combinations of:-

- |            |   |
|------------|---|
| 1 RYEGRASS | Provision of ryegrass:  |
| NONE       | None  |
| UNDRSOWN   | Undersown   |
| 2. APHIDS  | Aphids applied in April and early May:                                |
| NONE       | None  |
| METO FES   | Metopolophium festucae, on seven occasions, total 20 per square metre |

NOTE: Plots with ryegrass in 1982 were re-sown to ryegrass for 1983. Two extra plots were established for the 1983 combination RYEGRASS NONE, APHIDS NONE.

Basal applications: Manures: (5:14:30) at 340 kg. 'Nitro-Chalk' at 280 kg followed by 630 kg. Weedkillers: Methabenzthiazuron at 1.2 kg in 120 l. Dicamba, mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l with the prochloraz. Fungicides: Prochloraz at 0.40 kg. Carbendazim at 0.15 kg with maneb at 1.6 kg and tridemorph at 0.38 kg in 250 l.

Seed: Aquila: sown at 200 kg.  
Ryegrass: S24 sown at 22 kg.

Cultivations, etc.:- Ploughed: 31 Aug, 1982. NPK applied: 18 Sept. Spring-tine cultivated: 21 Sept. Spring-tine cultivated, wheat alone plots sown: 11 Oct. Wheat with undersown ryegrass sown: 12 Oct. Methabenzthiazuron applied: 16 Oct. First N applied: 15 Mar, 1983. Second N applied: 13 Apr. 'Herrisol' with prochloraz applied: 16 Apr. Carbendazim with maneb and tridemorph applied: 21 June. Combine harvested: 14 Aug.

NOTE: Cereal aphids were counted on seven occasions during June and July. Above-ground fauna were sampled on nine occasions between May and July. Ground predators were counted weekly from early May to early August. Parasites and fungal pathogens of aphids were assessed during June and July.

83/R/CS/271

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

APHIDS	NONE	METO FES	MEAN
RYEGRASS			
NONE	6.69	6.23	6.46
UNDRSOWN	5.43	5.35	5.39
MEAN	6.06	5.79	5.92

GRAIN MEAN DM% 87.3

PLOT AREA HARVESTED 0.01536