

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/WF/3 Wheat and Fallow - W. Wheat

Rothamsted Research

Rothamsted Research (1984) *83/R/WF/3 Wheat and Fallow - W. Wheat* ; Yields Of The Field Experiments 1983, pp 17 - 17 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/WF/3

WHEAT AND FALLOW

Object: To study the effects of fallowing on unmanured w. wheat -
Hoosfield.

The 128th year, w. wheat.

For previous years see 'Details' 1967, 1973 and 74-82/R/WF/3.

Whole plot dimensions: 9.60 x 211.

Treatments:

Each year there are two plots, one is sown to w. wheat, one is fallow; they alternate in successive years. The comparison of effects of three-year and one-year fallow, started in 1932, was made for the last time in 1982.

Standard applications:

Wheat plot: Weedkillers: Isoproturon at 2.1 kg and mecoprop, bromoxynil and ioxynil (as 'Brittox' at 3.5 l) in 250 l.

Seed: Flanders, dressed chlorfenvinphos, sown at 210 kg.

Cultivations, etc:-

Wheat plot: Ploughed: 7 Sept, 1982. Spring-tine cultivated: 29 Oct.
Heavy spring-tine cultivated, spring-tine cultivated, rotary harrowed, seed sown: 18 Jan, 1983. Weedkillers applied: 16 Apr.
Combine harvested: 10 Aug.

Fallow plot: Ploughed: 7 Sept, 1982. Heavy spring-tine cultivated: 25 May, 1983. Ploughed: 16 June. Spring-tine cultivated: 23 June.
Ploughed: 14 July. Spring-tine cultivated: 21 July.

GRAIN AND STRAW TONNES/HECTARE

	GRAIN	STRAW
YIELD	1.73	0.69
MEAN DM%	77.0	93.3
PLOT AREA HARVESTED	0.06009	