

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/AG/6 Agdell - W. Wheat

Rothamsted Research

Rothamsted Research (1984) *83/R/AG/6 Agdell - W. Wheat* ; Yields Of The Field Experiments 1983, pp 25 - 27 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/AG/6

AGDELL

Object: To study, by crop yields and soil analyses, the residual values of phosphate and potash applied in the period 1848-1951 and further dressings since 1964.

The 14th year of revised scheme, w. wheat.

For previous years see 'Details' 1967 and 1973, and 74-82/R/AG/6.

Treatments: All combinations of:-

Whole plots

1. OLDRESO Fertilizers and organic manures applied to roots every fourth year, in the period 1848-1948:

NONE	None
PKNAMG	P K Na Mg
NPKNAMGC	N P K Na Mg C

N: 48 kg N as sulphate of ammonia
P: 41 kg P as superphosphate
K: 224 kg K as sulphate of potash
Na: 16 kg Na as sulphate of soda
Mg: 11 kg Mg as sulphate of magnesia
C: Castor meal at 2240 kg supplying about 112 kg N

2. RN CROP Rotation 1848-1951 and crop in 1983:

FALLOW	With fallow: Roots (turnips or swedes), s. barley, fallow, w. wheat 1848-1951. Fallow in 1983
WHEAT	With legume: Roots, s. barley, legume (clover or beans), w. wheat 1848-1951. W. wheat 1983 (after w. beans 1982)

Half plots

3. 1964RESO Residues of 1964 treatments:

P
K

Quarter plots

4. PREVCROP Previous cropping 1958-69 on P-test half plots, 1958-70 on K-test half plots:

ARABLE	Arable or fallow
GRASS	Grass

83/R/AG/6

Sixteenth plots

5. P ₂ O ₅ 64	K ₂ O 64	Rates of 1964 treatments (kg):	
		P ₂ O ₅ to P-test half plots	K ₂ O to K-test half plots
0	0		
500	315		
1000	630		
2000	1260		

Thirty second plots

6.		To RN CROP WHEAT. Residues of P ₂ O ₅ applied 1970-72 (kg) and in 1980 and 1982 (kg):	
P ₂ O ₅ 722			
(0)0		None	
(375)300		375 total in 1970-72, 150 in 1980, 150 in 1982	
		To RN CROP WHEAT. Residues of K ₂ O applied 1973-76 (kg) and in 1980 and 1982 (kg):	
K ₂ O 762			
(0)0		None	
(870)600		870 total in 1973-76, 300 in 1980, 300 in 1982	

NOTE: Treatment combinations to thirty second plots of FALLOW plots are not shown above.

Standard applications:

W. wheat: Manures: 'Nitro-Chalk' at 130 kg followed by 770 kg.
Weedkillers: Mecoprop at 3.2 l with isoproturon at 2.1 kg in 250 l
applied with the prochloraz. Glyphosate at 1.4 kg in 250 l.
Fungicides: Prochloraz at 0.40 kg. Propiconazole at 0.12 kg in
250 l.

Seed: Avalon, sown at 180 kg.

Cultivations, etc.:-

W. wheat: Ploughed: 7 Sept, 1982. Spring-tine cultivated: 2 Oct.
Rotary harrowed, seed sown: 15 Oct. First N applied: 16 Mar, 1983.
Second N applied: 15 Apr. Mecoprop, isoproturon and prochloraz
applied: 16 Apr. Propiconazole applied: 17 June. Glyphosate
applied: 4 Aug. Combine harvested: 12 Aug.
Fallow: Ploughed: 7 Sept, 1982. Rotary cultivated: 24 May, 20 June,
1983. Spring-tine cultivated: 2 Aug.

83/R/AG/6

WHEAT P PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	OLDRES	NONE	PKNAMG		NPKNAMGC		
	P205 722	(0)0	(375)300	(0)0	(375)300	(0)0	(375)300
ARABLE	P205 64						
	0	7.57	9.21	7.87	7.99	7.55	8.06
	500	7.51	8.86	8.32	8.46	7.15	8.06
	1000	8.55	8.46	8.08	8.41	8.15	8.53
GRASS	2000	8.94	8.96	8.62	9.03	7.78	7.89
	0	5.99	9.35	6.96	8.56	6.13	7.62
	500	8.77	9.20	6.82	6.54	8.13	8.11
	1000	8.65	9.10	8.13	8.05	7.71	8.45
	2000	8.67	9.04	7.33	5.32	7.90	7.56

GRAIN MEAN DM% 86.3

WHEAT K PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	OLDRES	NONE	PKNAMG		NPKNAMGC		
	K20 762	(0)0	(870)600	(0)0	(870)600	(0)0	(870)600
ARABLE	K20 64						
	0	8.19	9.03	8.53	8.41	8.59	8.33
	315	8.26	9.58	8.13	8.32	7.92	7.33
	630	8.47	8.67	8.27	8.55	8.15	8.48
GRASS	1260	9.00	9.40	9.03	8.36	8.34	8.72
	0	8.50	8.98	7.99	8.45	8.11	8.55
	315	8.75	8.93	8.20	8.48	8.04	8.25
	630	8.71	9.12	8.91	8.61	8.18	8.77
	1260	8.63	9.43	8.36	9.07	8.36	8.77

GRAIN MEAN DM% 86.4

PLOT AREA HARVESTED (OLDRES NONE) 0.00131

PLOT AREA HARVESTED (REMAINDER) 0.00146