

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1983

[Full Table of Content](#)



## 83/R/B/4 and 83/W/B/4 Varieties and N - S. Barley

### Rothamsted Research

Rothamsted Research (1984) *83/R/B/4 and 83/W/B/4 Varieties and N - S. Barley* ; Yields Of The Field Experiments 1983, pp 257 - 259 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/B/4 and 83/W/B/4

SPRING BARLEY

VARIETIES AND N

Object: To study the yields of some of the newer varieties of s. barley at three rates of nitrogen - Rothamsted (R), Highfield IV and Woburn (W), Bull Field.

Sponsor: R. Moffitt.

Design: 2 randomised blocks of 3 plots split into 6.

Whole plot dimensions: 3.0 x 10.0.

Treatments: All combinations of:-

Whole plots

1. N Nitrogen fertilizer (kg N) as 'Nitro-Chalk', in addition to basals:

75  
113  
150

Sub plots

2. VARIETY Varieties:

ATEM  
CARNIVAL  
KYM  
PATTY  
TASMAN  
TRIUMPH

NOTE: At Woburn 75 kg N of the nitrogen treatments was applied to the seedbed as (20:10:10), the remainder and all at Rothamsted was applied as 'Nitro-Chalk'.

Basal applications:

Highfield IV (R): Manures: (0:18:36) at 690 kg, N at 65 kg as 'Nitro-Chalk'. Weedkillers: Dicamba with mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l. Fungicide: Prochloraz at 0.4 l in 250 l.

Bull Field (W): Manures: Magnesian limestone at 7.5 t, (20:10:10) at 377 kg. Weedkillers: Dicamba, with mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l. Fungicide: Tridemorph at 0.52 kg in 250 l.

Seed: Highfield IV (R) and Bull Field (W): Sown at 160 kg.

83/R/B/4 and 83/W/B/4

Cultivations, etc.:-

Highfield IV (R): PK applied: 18 Oct, 1982. Ploughed: 17 Dec. Spring-tine cultivated: 15 Mar, 1983. N treatments applied, spring-tine cultivated, seed sown: 17 Mar. Weedkillers applied: 24 May. Basal N applied: 27 May. Fungicide applied: 1 July. Combine harvested: 8 Aug. Previous crops: S. beans 1981, w. wheat 1982.  
 Bull Field (W): Magnesian limestone applied: 15 Sept, 1982. Ploughed: 15 Nov. Heavy spring-tine cultivated: 7 Mar, 1983. NPK applied: 10 Mar. Spring-tine cultivated with crumbler attached, seed sown: 11 Mar. Weedkillers applied: 26 May. Remaining N treatments applied: 3 June. Fungicide applied: 16 June. Combine harvested: 6 Aug. Previous crops: S. barley 1981 and 1982.

83/R/B/4 HIGHFIELD IV (R)

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	N	75	113	150	MEAN
VARIETY					
ATEM		6.81	6.11	7.10	6.67
CARNIVAL		6.63	5.53	7.06	6.41
KYM		7.37	6.45	7.37	7.06
PATTY		6.71	6.07	7.12	6.63
TASMAN		5.48	5.52	6.64	5.88
TRIUMPH		5.97	5.92	6.79	6.23
MEAN		6.49	5.94	7.01	6.48

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	VARIETY	N*
	VARIETY	
SED	0.284	0.493

\* FOR WITHIN SAME LEVEL OF N ONLY

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP.SP	15	0.493	7.6

GRAIN MEAN DM% 85.4

SUB PLOT AREA HARVESTED 0.00204

83/W/B/4 BULL FIELD (W)

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	N	75	113	150	MEAN
VARIETY					
ATEM		5.87	4.79	4.86	5.17
CARNIVAL		4.66	4.50	5.33	4.83
KYM		4.74	4.62	4.81	4.72
PATTY		4.63	4.38	3.61	4.21
TASMAN		3.62	4.09	3.33	3.68
TRIUMPH		4.49	4.28	4.26	4.34
MEAN		4.67	4.44	4.37	4.49

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	VARIETY	N*
	VARIETY	
-----		
SED	0.330	0.572

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP.SP	15	0.572	12.7

GRAIN MEAN DM% 86.1

SUB PLOT AREA HARVESTED 0.00275