

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/B/5 Sowing Dates and Insects - S. Barley

Rothamsted Research

Rothamsted Research (1984) *83/R/B/5 Sowing Dates and Insects - S. Barley*; Yields Of The Field Experiments 1983, pp 260 - 261 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/B/5

SPRING BARLEY

SOWING DATES AND INSECTS

Object: To study the effects of omethoate on insect pests and on yields of s. barley sown on two dates - Sawyers I.

Sponsor: G.C. Scott.

Design: 4 randomised blocks of 9 plots.

Whole plot dimensions: 9.0 x 10.0.

Treatments:

SDTE INS	Sowing dates and insecticides:
SE NONE	Sown 8 Mar, 1983, no insecticides
SE OME R	Sown 8 Mar, omethoate applied on 27 May, 13 June, 28 June, 12 July
SL NONE	Sown 14 Apr, no insecticides
SL OME 1	Sown 14 Apr, omethoate applied on 27 May
SL OME 2	Sown 14 Apr, omethoate applied on 13 June
SL OME 3	Sown 14 Apr, omethoate applied on 28 June
SL OME 4	Sown 14 Apr, omethoate applied on 12 July
SL OME R	Sown 14 Apr, omethoate applied 27 May, 13 June, 28 June, 12 July

NOTE: Omethoate was applied at 0.64 kg in 450 l.

Basal applications: Manures: 'Nitro-Chalk' at 500 kg followed by 250 kg. Weedkillers: Dicamba, mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l. Fungicide: Tridemorph at 0.52 kg in 250 l.

Seed: Triumph, dressed triadimenol plus fuberidazole, sown at 160 kg.

Cultivations, etc.: - Ploughed: 10 Nov, 1982. Spring-tine cultivated: 7 Mar, 1983. Spring-tine cultivated, early-sown plots rotary harrowed and sown: 8 Mar. Late-sown plots rotary harrowed and sown: 14 Apr. First N applied: 15 Apr. Weedkillers applied: 24 May. Second N applied: 26 May. Fungicide applied: 21 June. Combine harvested: 9 Aug. Previous crops: Potatoes 1981, s. barley 1982.

NOTES: (1) Aphids, thrips and stem borers were counted from the end of April to the middle of July.
(2) Components of yield were measured.

83/R/B/5

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

SDTE INS	
SE NONE	6.27
SE OME R	6.83
SL NONE	5.24
SL OME 1	4.89
SL OME 2	5.44
SL OME 3	5.85
SL OME 4	5.59
SL OME R	5.43
MEAN	5.64

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	SDTE INS
-----	-----
SED	0.424 MIN REP
	0.367 MAX-MIN

	SDTE INS
MAX-MIN	SL NONE V ANY OF REMAINDER
MIN REP	ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	25	0.600	10.6
GRAIN MEAN DM%	87.1		
PLOT AREA HARVESTED	0.00204		