Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1983



Full Table of Content

83/R/B/6 Mildew Sensitivity - S. Barley

Rothamsted Research

Rothamsted Research (1984) 83/R/B/6 Mildew Sensitivity - S. Barley; Yields Of The Field Experiments 1983, pp 264 - 265 - DOI: https://doi.org/10.23637/ERADOC-1-44

83/R/B/6

SPRING BARLEY

MILDEW SENSITIVITY

Object: To study the effects of varieties with differing resistance genes on the sensitivity of powdery mildew (Erysiphe graminis) to fungicides - Fosters Corner.

Sponsor: D.W. Hollomon.

Design: 2 randomised blocks of 12 plots.

Whole plot dimensions: 9.0×9.0 .

Treatments: All combinations of:-

1. VARIETY Varieties:

CARNIVAL TRIUMPH

2. FUNG SD Fungicidal seed dressings:

NONE None

ETHIRIMO Ethirimol at 4 g per kg seed TR DMNOL Triadimenol at 0.4 g per kg seed

FUNG SP Fungicidal foliar spray:

NONE None

TR DMFON Triadimefon at 0.12 kg in 250 l on 17 June, 1983

NOTE: The seed was sown at 160 kg.

Basal applications: Manures: (0:18:36) at 690 kg. 'Nitro-Chalk' at 450 kg followed by 250 kg. Weedkillers: Dicamba, mecoprop and MCPA (as 'Herrisol' at 5.0 1) in 250 1.

Cultivations, etc.:- PK applied: 19 Oct, 1982. Ploughed: 25 Oct. Springtine cultivated: 8 Mar, 1983. First N applied: 10 Mar. Spring-tine cultivated: 11 Mar. Seed sown: 12 Mar. Weedkillers applied: 24 May. Second N applied: 26 May. Combine harvested: 8 Aug. Previous crops: S. beans 1981, w. wheat 1982.

NOTE: Mildew was assessed on six occasions during May, June and July; scald (Rhynchosporium secalis) was also recorded in May and June.

Sensitivity of mildew to triadimenol was assessed by bioassay on two occasions.

83/R/B/6

GRAIN TONNES/HECTARE

**** TABLES OF MEANS ****

FUNG SD	NONE	ETHIRIMO	TR DMNOL	MEAN		
VARIETY	6 40	6.51	6.98	6.66		
CARNIVAL	6.49					
TRIUMPH	6.00	6.01	5.95	5.99		
MEAN	6.25	6.26	6.47	6.33		
FUNG SP VARIETY	NONE	TR DMFON	MEAN			
CARNIVAL	6.20	7.13	6.66			
TRIUMPH	5.28	6.70	5.99			
MEAN	5.74	6.91	6.33			
FUNG SP	NONE	TR DMFON	MEAN.			
NONE	5.68	6.81	6.25			
	5.72	6.80				
ETHIRIMO						
TR DMNOL	5.81	7.13	6.47			
MEAN	5.74	6.91	6.33			
FUNG SD	NONE		ETHIRIMO	TR	DMNOL	
		TR DMFON	NONE TR	DMEON	NONE TR	DMFON
FUNG SP	NUNE	IK DITON	NONE IN	Drii Oil	HONE II	C DI'II OII
VARIETY						7.56
CARNIVAL	6.11	6.87		6.95	6.41	7.56
TRIUMPH	5.26	6.74	5.37	6.66	5.22	6.69

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	VARIETY	FUNG SD	FUNG SP	VARIETY FUNG SD
SED	0.196	0.240	0.196	0.339
TABLE	VARIETY FUNG SP	FUNG SD FUNG SP	VARIETY FUNG SD FUNG SP	
SED	0.277	0.339	0.480	

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV%

BLOCK.WP 11 0.480 7.6

GRAIN MEAN DM% 87.2

PLOT AREA HARVESTED 0.00248