

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1983

[Full Table of Content](#)



## 83/R/BE/13 Times of Applying Erynia - S. Beans

### Rothamsted Research

Rothamsted Research (1984) *83/R/BE/13 Times of Applying Erynia - S. Beans* ; Yields Of The Field Experiments 1983, pp 294 - 295 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/BE/13

SPRING BEANS

TIMES OF APPLYING ERYNIA

Object: To study the effects of rates and times of applying the aphid-pathogenic fungus *Erynia neoaphidis* on the numbers of black aphids (*Aphis fabae*) and on the yield of s. beans - W. Barnfield II.

Sponsor: N. Wilding.

Design: 4 randomised blocks of 7 plots.

Whole plot dimensions: 2.67 x 2.13.

Treatments:

APH CONT            Chemical and biological aphid control:

NONE                None

                      Pirimicarb at 0.14 kg in 340 l:

PIRIM E            Applied on 10 June, 1983

PIRIM L            Applied on 29 June

*Erynia neoaphidis* applied as a powder of mummified aphids:

E NEO1 E           At 0.5 mg per plant on 9 June

E NEO1 L           At 0.5 mg per plant on 28 June

E NEO2 E           At 5.0 mg per plant on 9 June

E NEO2 L           At 5.0 mg per plant on 28 June

NOTE: Basal irrigation was applied as follows (mm water):

21 June	20
28 June	25
5 July	12.5
12 July	12.5
15 July	12.5
23 July	12.5
Total	95 mm

Basal applications: Manures: Chalk at 5.0 t. Weedkillers: Glyphosate at 1.4 kg in 250 l. Trietazine with simazine (as 'Aventox' at 2.4 l) in 250 l.

Seed: Minden, sown at 230 kg.

Cultivations, etc.: - Chalk applied: 16 Sept, 1982. Glyphosate applied: 19 Oct. Ploughed: 25 Nov. Spring-tine cultivated twice: 8 Mar, 1983. Seed sown: 9 Mar. Rolled, trietazine and simazine applied: 12 Mar. Combine harvested: 12 Aug. Previous crops: S. barley 1981, w. wheat 1982.

NOTES: (1) Aphid numbers were estimated weekly during June and July.  
(2) Samples of live aphids were examined for infection with *Erynia* and other fungal pathogens.

83/R/BE/13

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

APH CONT	
NONE	4.75
PIRIM E	5.34
PIRIM L	4.89
E NEO1 E	4.32
E NEO1 L	4.33
E NEO2 E	4.72
E NEO2 L	4.69
MEAN	4.72

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	APH CONT
-----	-----
SED	0.203

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	18	0.287	6.1

GRAIN MEAN DM% 91.3

PLOT AREA HARVESTED 0.00024