

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/BE/15 Control of Rust - S. Beans

Rothamsted Research

Rothamsted Research (1984) *83/R/BE/15 Control of Rust - S. Beans* ; Yields Of The Field Experiments 1983, pp 298 - 299 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/BE/15

SPRING BEANS

CONTROL OF RUST

Object: To study the effects of fungicides on the control of rust (*Uromyces fabae*) and on the yield of spring beans - Long Hoos IV 5.

Sponsors: D.H. Lapwood, J. McEwen, D.P. Yeoman.

Design: 3 randomised blocks of 12 plots.

Whole plot dimensions: 2.03 x 2.13.

Treatments: All combinations of:-

1. C S FUNG Fungicide to control chocolate spot but not rust:
 NONE None
 BENOMYL Benomyl at 0.50 kg in 340 l on 7 July, 1983
2. RUSTFUNG Fungicides to control rust:
 MAN+MANC Maneb at 0.8 kg + mancozeb at 0.8 kg in 340 l
 PROPICON Propiconazole at 0.12 kg in 340 l
3. RFNGTIME Times of applying fungicides to control rust:
 ONCE Once on 18 July
 TWICE Twice, on 18 and 27 July

plus extra treatments:

EXTRA

- | | |
|---------|--|
| NONE | None (duplicated) |
| BENOMYL | Benomyl at 0.50 kg in 340 l on 7 July (duplicated) |

Basal applications: Manures: Chalk at 2.9 t. Muriate of potash at 520 kg.
Weedkillers: Trietazine at 1.1 kg with simazine at 0.16 kg in 220 l.
Insecticides: Pirimicarb at 0.14 kg in 340 l on two occasions;
permethrin at 0.15 kg in 340 l on two occasions.

Seed: Minden, sown at 280 kg.

Cultivations, etc.: - Muriate of potash applied: 16 Sept, 1982. Chalk applied: 30 Sept. Ploughed: 11 Oct. Spring-tine cultivated, seed sown: 8 Mar, 1983. Weedkillers applied: 17 Mar. Permethrin applied: 4 May, 20 May. Pirimicarb applied; 13 June, 13 July. Harvested by hand: 16 Aug. Previous crops: Potatoes 1981, s. barley 1982.

NOTE: Plant counts were made after establishment. The incidence of chocolate spot and rust were assessed from early July until maturity. Components of yield were measured at maturity.

83/R/BE/15

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

RUSTFUNG	MAN+MANC	PROPICON	MEAN
C S FUNG			
NONE	3.68	4.22	3.95
BENOMYL	3.86	4.39	4.13
MEAN	3.77	4.30	4.04
RFNGTIME	ONCE	TWICE	MEAN
C S FUNG			
NONE	4.31	3.59	3.95
BENOMYL	3.83	4.42	4.13
MEAN	4.07	4.00	4.04
RFNGTIME	ONCE	TWICE	MEAN
RUSTFUNG			
MAN+MANC	3.93	3.62	3.77
PROPICON	4.21	4.39	4.30
MEAN	4.07	4.00	4.04
RUSTFUNG	MAN+MANC	PROPICON	
RFNGTIME	ONCE	TWICE	ONCE
C S FUNG			TWICE
NONE	4.20	3.15	4.42
BENOMYL	3.65	4.08	4.01
EXTRA	NONE	BENOMYL	MEAN
	3.64	3.51	3.58
GRAND MEAN	3.88		

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	C S FUNG	RUSTFUNG	RFNGTIME
SED	0.253	0.179	0.179	0.179
TABLE	C S FUNG	C S FUNG	RUSTFUNG	C S FUNG
	RUSTFUNG	RFNGTIME	RFNGTIME	RUSTFUNG
				RFNGTIME
SED	0.253	0.253	0.253	0.358

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	24	0.439	11.3
GRAIN MEAN DM%	89.3		
PLOT AREA HARVESTED	0.00015		