

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1983

[Full Table of Content](#)



## 83/R/SW/1 Electrostatic Spraying and Mildew - Swedes

### Rothamsted Research

Rothamsted Research (1984) *83/R/SW/1 Electrostatic Spraying and Mildew - Swedes* ; Yields Of The Field Experiments 1983, pp 318 - 319 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/R/SW/1

SWEDES

ELECTROSTATIC SPRAYING AND MILDEW

Object: To compare the effects of electrostatic and conventional sprayers on disease control and on yield of swedes - Pennell's Piece.

Sponsors: A.J. Arnold, G.R. Cayley, P. Etheridge, D.C. Griffiths, F.T. Phillips, B.J. Pye, G.C. Scott.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 3.05 x 13.7.

Treatments: All combinations of:-

1. SPRAYER                      Spraying machines:

CNVNTIAL	Conventional
ELECT NA	New electrostatic A
ELECT NB	New electrostatic B
ELECT O	Older electrostatic

2. NUARIMOL                      Rates of applying nuarimol fungicide(g):

40  
80

plus one extra treatment not sprayed

EXTRA

NONE                      None (duplicated)

- NOTES: (1) The new electrostatic A sprayer had spinning-disc nozzles and applied spray in 6.95 l.  
(2) The new electrostatic B and older electrostatic sprayers had spinning-cone nozzles, applied spray in 10.6 l, the former was mounted at an angle of 30° to the vertical, the latter vertical.  
(3) Nuarimol was applied on 11 Aug, 1983 by all sprayers, in 350 l by the conventional sprayer.

Basal applications: Manures: Chalk at 5.0 t. (0:18:36) at 630 kg. 'Nitro-Chalk' at 380 kg. Weedkillers: Glyphosate at 1.4 kg in 250 l. Paraquat at 0.98 kg ion in 900 l. Trifluralin at 1.1 kg in 250 l. Insecticide: Pirimicarb at 0.075 kg to all treatments, applied in 380 l by conventional sprays to all treatments except one of the duplicate EXTRA : NONE treatments, by electrostatic sprayer in 8.3 l.

Seed: Doon Major, dressed gamma HCH with captan, sown at 2.0 kg.

83/R/SW/1

Cultivations, etc.:- Glyphosate applied: 28 Oct, 1982. Chalk applied: 29 Nov. Ploughed: 8 Dec. N and PK applied: 28 Apr, 1983. Paraquat applied: 14 May. Heavy spring-tine cultivated, trifluralin applied, rotary harrowed: 23 May. Seed sown: 25 May. Plants singled: 29 June. Insecticide applied: 25 July. Harvested: 10 Nov. Previous crops: Peas 1981, s. barley 1982.

NOTE: Samples for chemical analysis were taken immediately after spraying. Mildew assessments were made at intervals throughout the season.

TOTAL TUBERS TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

SPRAYER	CNVNTIAL	ELECT NA	ELECT NB	ELECT O	MEAN
NUARIMOL					
40	29.2	32.7	36.0	35.1	33.2
80	33.7	34.5	33.2	32.3	33.4
MEAN	31.5	33.6	34.6	33.7	33.3
NUARIMOL NONE	23.9				
GRAND MEAN	31.4				

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	NUARIMOL	SPRAYER	NUARIMOL SPRAYER
-----			
SED	3.40	4.80	6.79

SED FOR COMPARING NUARIMOL NONE WITH A MEAN IN NUARIMOL.SPRAYER TABLE IS 5.88

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	19	8.32	26.5
PLOT AREA HARVESTED	0.00084		