

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/W/G/1 Fluid Drilling and Nematicides - Grass

Rothamsted Research

Rothamsted Research (1984) *83/W/G/1 Fluid Drilling and Nematicides - Grass* ; Yields Of The Field Experiments 1983, pp 320 - 322 - DOI: <https://doi.org/10.23637/ERADOC-1-44>

83/W/G/1

GRASS

FLUID DRILLING AND NEMATICIDES

Object: To study the effects of rates of oxamyl incorporated in a fluid gel at sowing on the establishment and yield of ryegrass - Woburn Butt Close.

Sponsor: A.M. Spaul.

Design: 5 randomised blocks of 5 plots.

Whole plot dimensions: 1.22 x 3.05.

Treatments:

OXAMYL	Rates of oxamyl (kg) incorporated in a fluid gel at sowing:
--------	-------------------------------------------------------------

0.0	
0.2	
0.5	
2.0	
5.0	

Basal applications: Manures: (10:10:15+4.5 Mg) at 750 kg, N at 75 kg as 'Nitro-Chalk'. Weedkillers: Glyphosate at 1.5 kg in 280 l, mecoprop with bromoxynil and ioxynil (as 'Brittox' at 2.5 l) in 280 l.

Seed: RVP Italian ryegrass, sown at 20 kg.

Cultivations, etc.:- Glyphosate applied: 6 Oct, 1982. Ploughed: 2 Nov. Spring-tine cultivated: 15 Mar, 1983. NPK + Mg applied: 13 Apr. Treatments applied, seeds sown: 15 Apr. Hand weeded: 17 June. 'Brittox' applied: 22 June. Cut twice: 1 Aug, 12 Oct. N applied: 1 Aug. Previous crops: Potatoes 1981, w. wheat 1982.

NOTE: Nematodes were assessed in soil samples taken in April before sowing and in October after the second cut.

83/W/G/1

1ST CUT (1/8/83) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

OXAMYL	0.0	0.2	0.5	2.0	5.0	MEAN
	2.33	2.00	1.81	2.29	1.88	2.06

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	OXAMYL
SED	0.311

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	16	0.491	23.8

1ST CUT MEAN DM% 34.3

2ND CUT (12/10/83) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

OXAMYL	0.0	0.2	0.5	2.0	5.0	MEAN
	1.88	1.50	1.72	1.87	1.58	1.71

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	OXAMYL
SED	0.274

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	16	0.433	25.3

2ND CUT MEAN DM% 16.5

83/W/G/1

TOTAL OF 2 CUTS DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

OXAMYL	0.0	0.2	0.5	2.0	5.0	MEAN
	4.21	3.50	3.53	4.15	3.46	3.77

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	OXAMYL
-----	-----
SED	0.421

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	16	0.665	17.6

TOTAL OF 2 CUTS MEAN DM% 25.4

PLOT AREA HARVESTED 0.00025