

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1983

[Full Table of Content](#)



83/R/RN/17 Rates of P and K to the Subsoil - S. Beans, W. Wheat, Potatoes, S. Barley

Rothamsted Research

Rothamsted Research (1984) *83/R/RN/17 Rates of P and K to the Subsoil - S. Beans, W. Wheat, Potatoes, S. Barley*; Yields Of The Field Experiments 1983, pp 89 - 95 - DOI:

<https://doi.org/10.23637/ERADOC-1-44>

83/R/RN/17

RATES OF P AND K TO THE SUBSOIL

Object: To study the effects of a range of rates and frequencies of application of P and K to the subsoil, singly and together, on the yields and nutrient uptakes of a rotation of crops - Meadow.

Sponsors: J. McEwen, A.E. Johnston.

The third year, potatoes, s. barley, s. beans, w. wheat.

For previous years see 81-82/R/RN/17.

Design: 4 series (for crops) each of 40 plots.

Whole plot dimensions: 3.0 x 14.0.

Treatments to each series:

TREATMNT Extra P and K and primary cultivation tool in autumn 1980 only except on R plots, treatments repeated each autumn:

	P ₂ O ₅ (kg)	K ₂ O(kg)	Tool	
- - -	0	0	Plough	(duplicated)
P6 K6 T	1000	500 to topsoil	"	(")
- - S	0	0 " "	Wye double-digger	(four plots)
- - SR	0	0 " "	" "	(duplicated)
P2 - SR	63	0 to subsoil	" "	"
P3 - S	125	0 " "	" "	"
P4 - S	250	0 " "	" "	"
P5 - S	500	0 " "	" "	(duplicated)
P6 - S	1000	0 " "	" "	"
- K2 SR	0	31 " "	" "	"
- K3 S	0	63 " "	" "	"
- K4 S	0	125 " "	" "	"
- K5 S	0	250 " "	" "	(duplicated)
- K6 S	0	350 " "	" "	"
P1 K1 SR	31	16 " "	" "	"
P1 K3 SR	31	63 " "	" "	"
P2 K2 SR	63	31 " "	" "	"
P3 K1 SR	125	16 " "	" "	"
P3 K3 SR	125	63 " "	" "	"
P3 K4 S	125	125 " "	" "	"
P4 K3 S	250	63 " "	" "	"
P4 K4 S	250	125 " "	" "	"
P4 K5 S	250	250 " "	" "	(duplicated)
P4 K6 S	250	350 " "	" "	"
P5 K4 S	500	125 " "	" "	(duplicated)
P5 K5 S	500	250 " "	" "	"
P5 K6 S	500	350 " "	" "	"
P6 K4 S	1000	125 " "	" "	"
P6 K5 S	1000	250 " "	" "	"
P6 K6 S	1000	350 " "	" "	"

83/R/RN/17

- NOTES: (1) Subsoiling was done with the Wye double-digger which turns a furrow with a conventional plough share, to a depth of 23 cm, and at the same time rotary cultivates the bottom of the adjacent furrow to a further depth of 15 cm. When applying P and K this was distributed ahead of the rotary cultivator.
- (2) The topsoil PK dressing was equally divided before and after ploughing.
- (3) All plots other than R were conventionally ploughed in autumn 1981 and 1982.
- (4) The rate of 350 kg K₂O applied was in error for 500 kg K₂O.

Standard applications:

- Potatoes: Manures: (10:10:15 + 4.5 Mg) at 1960 kg. Weedkillers: Paraquat at 0.56 kg ion with linuron at 1.0 l in 500 l. Fungicides: Mancozeb at 1.4 kg in 250 l on three occasions. Fentin hydroxide at 0.28 kg in 250 l on five occasions, with the insecticide on the first four. Insecticide: Pirimicarb at 0.14 kg. Haulm desiccant: BOV at 170 l.
- S. barley: Manures: (20:10:10) at 450 kg. 'Nitro-Chalk' at 250 kg. Weedkillers: Dicamba, mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l. Fungicide: Prochloraz at 0.40 kg in 250 l.
- S. beans: Weedkillers: Trietazine and simazine (as 'Aventox' at 2.4 l) in 250 l. Insecticides: Pirimicarb at 0.14 kg in 250 l. Phorate at 5.6 kg.
- W. wheat: Manures: (0:18:36) at 350 kg combine drilled. 'Nitro-Chalk' at 750 kg. Weedkillers: Dicamba, mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l. Fungicide: Propiconazole at 0.12 kg in 250 l.

Seed: Potatoes: Pentland Crown.

- S. barley: Triumph, seed dressed with ethirimol, sown at 160 kg.
- S. beans: Minden, sown at 280 kg.
- W. wheat: Avalon, sown at 210 kg.

Cultivations, etc.:-

All crops: Treatments applied by double digger: 15-17 Nov, 1982.
Ploughed: 18 Nov.

- Potatoes: Spring-tine cultivated twice: 7 Mar, 1983. NPK Mg applied: 28 Apr. Spiked rotary cultivated, potatoes planted: 29 Apr. Rotary ridged: 6 May. Weedkillers applied: 24 May. Mancozeb applied: 22 June. Fentin hydroxide with insecticide applied: 1 July, 8 July, 18 July and 28 July. Fentin hydroxide applied: 11 Aug. Mancozeb applied: 25 Aug, 9 Sept. Haulm mechanically destroyed: 11 Oct. BOV applied: 19 Oct. Lifted: 27 Oct.
- S. barley: Spring-tine cultivated, NPK applied, spring-tine cultivated a second time: 7 Mar, 1983. Rotary harrowed, seed sown: 8 Mar. 'Herrisol' applied: 23 May. N applied: 26 May. Fungicide applied: 23 June. Combine harvested: 3 Aug.
- S. beans: Spring-tine cultivated, phorate applied, spring-tine cultivated a second time, rotary harrowed, seed sown: 7 Mar, 1983. Weedkillers applied: 12 Mar. Pirimicarb applied: 16 June. Combine harvested: 12 Aug.
- W. wheat: Heavy spring-tine cultivated twice, spring-tine cultivated twice, seed sown: 18 Jan, 1983. N applied: 16 Apr. Weedkillers applied: 28 Apr. Fungicide applied: 17 June. Combine harvested: 12 Aug.

83/R/RN/17

SERIES III POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	43.6
P6 K6 T	45.4
- - S	41.5
- - SR	38.8
P2 - SR	36.1
P3 - S	45.8
P4 - S	45.3
P5 - S	42.1
P6 - S	46.4
- K2 SR	39.8
- K3 S	44.6
- K4 S	45.2
- K5 S	41.8
- K6 S	46.5
P1 K1 SR	39.2
P1 K3 SR	45.2
P2 K2 SR	36.4
P3 K1 SR	43.1
P3 K3 SR	39.3
P3 K4 S	41.0
P4 K3 S	42.4
P4 K4 S	39.7
P4 K5 S	44.2
P4 K6 S	46.9
P5 K4 S	43.1
P5 K5 S	43.9
P5 K6 S	47.6
P6 K4 S	47.5
P6 K5 S	45.3
P6 K6 S	43.5
MEAN	42.9

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	4.73 MIN REP
	3.74 MAX-MIN

* SED APPLIES ONLY TO - - -, P6 K6 T, - - S, - - SR, P5 - S, - K5 S, P4 K5 S AND P5 K4 S

TREATMNT
 MAX-MIN - - S V ANY OF REMAINDER
 MIN REP ANY OF THE REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	3.34	7.8

83/R/RN/17

SERIES III POTATOES

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

TREATMNT	
- - -	93.8
P6 K6 T	94.7
- - S	93.5
- - SR	95.2
P2 - SR	93.9
P3 - S	93.5
P4 - S	93.4
P5 - S	94.1
P6 - S	96.2
- K2 SR	92.6
- K3 S	94.6
- K4 S	93.9
- K5 S	95.1
- K6 S	95.5
P1 K1 SR	93.5
P1 K3 SR	95.9
P2 K2 SR	93.7
P3 K1 SR	95.2
P3 K3 SR	93.3
P3 K4 S	95.5
P4 K3 S	94.8
P4 K4 S	93.7
P4 K5 S	95.3
P4 K6 S	96.9
P5 K4 S	94.1
P5 K5 S	94.7
P5 K6 S	95.0
P6 K4 S	94.5
P6 K5 S	94.4
P6 K6 S	93.6
MEAN	94.4

PLOT AREA HARVESTED 0.00210

83/R/RN/17

SERIES IV SPRING BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	7.30
P6 K6 T	7.43
- - S	6.82
- - SR	6.42
P2 - SR	6.35
P3 - S	6.66
P4 - S	6.98
P5 - S	7.09
P6 - S	7.05
- K2 SR	6.87
- K3 S	7.24
- K4 S	7.37
- K5 S	6.94
- K6 S	7.56
P1 K1 SR	6.81
P1 K3 SR	7.33
P2 K2 SR	5.68
P3 K1 SR	7.13
P3 K3 SR	6.77
P3 K4 S	7.51
P4 K3 S	7.42
P4 K4 S	7.37
P4 K5 S	6.93
P4 K6 S	7.39
P5 K4 S	6.98
P5 K5 S	7.24
P5 K6 S	7.10
P6 K4 S	7.51
P6 K5 S	7.40
P6 K6 S	6.34
MEAN	7.01

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.336 MIN REP
	0.266 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.238	3.4
GRAIN MEAN DM%	85.7		
PLOT AREA HARVESTED	0.00286		

83/R/RN/17

SERIES I SPRING BEANS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	4.17
P6 K6 T	4.49
- - S	4.32
- - SR	4.61
P2 - SR	4.72
P3 - S	4.82
P4 - S	4.48
P5 - S	4.44
P6 - S	4.46
- K2 SR	4.34
- K3 S	4.28
- K4 S	4.01
- K5 S	4.28
- K6 S	4.41
P1 K1 SR	4.53
P1 K3 SR	4.24
P2 K2 SR	4.13
P3 K1 SR	4.34
P3 K3 SR	4.69
P3 K4 S	4.16
P4 K3 S	4.39
P4 K4 S	4.73
P4 K5 S	4.37
P4 K6 S	4.08
P5 K4 S	4.66
P5 K5 S	4.44
P5 K6 S	4.82
P6 K4 S	4.39
P6 K5 S	4.58
P6 K6 S	4.57
MEAN	4.42

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.324 MIN REP
	0.257 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.229	5.2

GRAIN MEAN DM% 85.5

PLOT AREA HARVESTED 0.00386

83/R/RN/17

SERIES II WINTER WHEAT

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	6.52
P6 K6 T	7.40
- - S	6.81
- - SR	6.06
P2 - SR	6.45
P3 - S	6.72
P4 - S	6.98
P5 - S	7.06
P6 - S	6.70
- K2 SR	5.65
- K3 S	7.04
- K4 S	6.78
- K5 S	6.44
- K6 S	5.42
P1 K1 SR	5.62
P1 K3 SR	6.14
P2 K2 SR	6.04
P3 K1 SR	6.54
P3 K3 SR	6.51
P3 K4 S	6.61
P4 K3 S	7.23
P4 K4 S	7.33
P4 K5 S	6.66
P4 K6 S	6.69
P5 K4 S	6.76
P5 K5 S	7.16
P5 K6 S	6.44
P6 K4 S	7.51
P6 K5 S	5.57
P6 K6 S	7.41
MEAN	6.64

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.362 MIN REP
	0.286 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.256	3.9

GRAIN MEAN DM% 85.7

PLOT AREA HARVESTED 0.00286