Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1984



Full Table of Content

# 84/R/CS/133 Control of Pathogens - Maize

## **Rothamsted Research**

Rothamsted Research (1985) 84/R/CS/133 Control of Pathogens - Maize; Yields Of The Field Experiments 1984, pp 127 - 128 - DOI: https://doi.org/10.23637/ERADOC-1-32

### 84/R/CS/133

#### CONTROL OF PATHOGENS

Object: To study the effects of a range of chemicals on incidence of pathogens and yield of continuous maize - Long Hoos VI/VII 6.

Sponsors: A.J. Barnard, K.E. Fletcher, D.J. Hooper, D. Hornby, R.T. Plumb.

The 11th year, forage maize.

For previous years see 74-83/R/CS/133.

Design: 3 randomised blocks of 9 plots.

Whole plot dimensions: 2.13 x 18.3.

Treatments:-

CHEMICAL Chemicals applied annually except as stated:

NONE None (2 plots per block)

ALDICARB Aldicarb, 4.5 kg as granules to seedbed BENOMYL Benomyl, 11.2 kg as dust to seedbed

DAZOMET Dazomet, 450 kg as granules in early spring (not applied

1975, 1979 and 1981)

PERMETH Permethrin, as foliar spray (0.15 kg in 1979, 0.05 kg in

1984)

PHORATE Phorate, 1.68 kg as granules to seedbed

PIRIMICA Pirimicarb, 0.14 kg as foliar spray (1979 & 1984 only) Benomyl + dazomet (not applied 1975, 1979 & 1981)

+ phorate, at above rates and times

NOTE: Permethrin and pirimicarb were applied in 340 l in 1979, 220 l in 1984.

Basal applications: Manures: 'Nitro-Chalk' at 660 kg. Weedkiller: Atrazine at 1.7 l in 220 l.

Seed: Fronica, sown at 100,000 seeds per hectare.

Cultivations, etc.:- Ploughed: 3 Nov, 1983. Spring-tine cultivated dazomet plots only: 21 Mar, 1984. Dazomet applied and these plots only rotary cultivated: 22 Mar. Spring-tine cultivated: 10 May. Remaining seedbed treatments applied, power harrowed, seed sown: 11 May. Weedkiller applied: 14 May. N applied: 18 May. Foliar treatments applied: 17 July. Harvested by hand: 30 Oct.

84/R/CS/133

FORAGE MAIZE DRY MATTER TONNES/HECTARE

\*\*\*\* TABLES OF MEANS \*\*\*\*

CHEMICAL 11.62 NONE ALDICARB 11.53 BENOMYL 11.26 12.81 DAZOMET PERMETH 11.71 PHORATE 11.71 11.66 PIRIMICA BE+DA+PH 15.33 MEAN 12.14

\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*

TABLE CHEMICAL 0.959 MIN REP SED 0.831 MAX-MIN

CHEMICAL

MAX-MIN NONE V ANY OF REMAINDER MIN REP ANY OF REMAINDER

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM DF SE CV% BLOCK. WP 17 1.175 9.7

FORAGE MEAN DM% 33.6

PLOT AREA HARVESTED 0.00059