

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/CS/247 Organic Matter and Earthworm Inoculation - S. Barley

Rothamsted Research

Rothamsted Research (1985) *84/R/CS/247 Organic Matter and Earthworm Inoculation - S. Barley* ; Yields Of The Field Experiments 1984, pp 152 - 153 - DOI:

<https://doi.org/10.23637/ERADOC-1-32>

84/R/CS/247

ORGANIC MATTER AND EARTHWORM INOCULATION

Object: To study methods of inoculating earthworms into arable soil and the influence of organic materials on subsequent multiplication and spread - Hoosfield.

Sponsor: C.A. Edwards.

The fifth year, s. barley.

For previous years see 80-83/R/CS/247.

Design: 3 randomised blocks of 9 plots.

Whole plot dimensions: 7.85 x 7.62.

Treatments: All combinations of:-

1. WORMINOC(80) Earthworms and inoculation method for 1980 crop only:

NONE	None
	Earthworms (<i>Lumbricus terrestris</i>) applied at 16,700 per hectare in November 1979:
EVEN	Evenly spaced throughout
CONC	Concentrated in metre squares, 100 earthworms per square metre

2. ORG MATT Forms of organic matter:

NONE	None
STR	Straw at 6.50 t for 1980, 3.25 t for 1981 and 1982
STR+FYM	Straw at 6.50 t for 1980, 3.25 t for 1981 and 1982 plus farmyard manure at 40 t for each year including 1984

Basal applications: Manures: (20:10:10) at 630 kg. Weedkillers: Paraquat at 0.4 kg ion in 250 l. 3, 6-dichloropicolinic acid at 0.05 kg and bromoxynil at 0.24 kg with mecoprop (as 'CMPP' at 3.0 l) in 250 l with the fungicide. Fungicide: Tridemorph at 0.52 kg.

Seed: Triumph, seed dressed with ethirimol, sown at 160 kg.

Cultivations, etc.: - Paraquat applied: 21 Oct, 1983. Deep spring-tine cultivated twice: 11 Nov. FYM treatment applied: 24 Nov. NPK applied: 8 Mar, 1984. Spring-tine cultivated twice, seed sown: 10 Mar. 3, 6-dichloropicolinic acid, bromoxynil, mecoprop and tridemorph applied: 16 May. Combine harvested: 17 Aug.

NOTE: Soil fauna were estimated from soil cores taken monthly from April to August and from pitfall trapping in the same period. Earthworm samples were taken in the autumn.

84/R/CS/247

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

ORG MATT WORMINOC(80)	NONE	STR	STR+FYM	MEAN
NONE	6.79	6.73	7.34	6.95
EVEN	6.77	6.39	6.55	6.57
CONC	6.52	6.83	6.65	6.67
MEAN	6.69	6.65	6.85	6.73

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	WORMINOC(80)	ORG MATT	WORMINOC(80) ORG MATT
SED	0.269	0.269	0.466

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	16	0.571	8.5

GRAIN MEAN DM% 84.7

PLOT AREA HARVESTED 0.00244