

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/CS/279 Nematicides and Stem Nematode - Lucerne

Rothamsted Research

Rothamsted Research (1985) *84/R/CS/279 Nematicides and Stem Nematode - Lucerne* ; Yields Of The Field Experiments 1984, pp 161 - 169 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/CS/279

NEMATICIDES AND STEM NEMATODE

Object: To study, on sites initially free from or infested by stem nematode (*Ditylenchus dipsaci*), the effects of nematicides on lucerne - Long Hoos V 5 (healthy) and Long Hoos IV 2 (infested).

Sponsor: A.G. Whitehead.

The third year, lucerne.

For previous years see 82-83/R/CS/279.

Design: On each site: 3 randomised blocks of 14 plots.

Whole plot dimensions: 1.2 x 3.7.

Treatments (applied to HEALTHY and INFESTED sites):

| TREATMNT | Varieties, rates and methods of applying nematicides: |
|----------|---|
| V 0 | Vertus, untreated |
| V A1 | Vertus, aldicarb at 1.5 kg in seed furrows in 1982 |
| E 0 | Europe, untreated |
| E A1 | Europe, aldicarb at 1.5 kg in seed furrows in 1982 |
| E A2 | Europe, aldicarb at 3.0 kg in seed furrows in 1982 |
| E A1 A1 | Europe, aldicarb at 1.5 kg in seed furrows in 1982, repeated after each cut in 1982 and in spring and after each cut thereafter |
| E A1 T1 | Europe, aldicarb at 1.5 kg in seed furrows in 1982, thiabendazole at 1.5 kg over the rows in spring 1983 and 1984 |
| E A2 T2 | Europe, aldicarb at 3.0 kg in seed furrows in 1982, thiabendazole at 3.0 kg over the rows in spring 1983 and 1984 |
| E C1 | Europe, carbofuran at 1.5 kg in seed furrows in 1982 |
| E C2 | Europe, carbofuran at 3.0 kg in seed furrows in 1982 |
| E C1 T1 | Europe, carbofuran at 1.5 kg in seed furrows in 1982, thiabendazole at 1.5 kg over the rows in spring 1983 and 1984 |
| E C2 T2 | Europe, carbofuran at 3.0 kg in seed furrows in 1982, thiabendazole at 3.0 kg over the rows in spring 1983 and 1984 |
| E T1 T1 | Europe, thiabendazole at 1.5 kg over the rows at sowing in 1982 and in spring 1983 and 1984 |
| E T2 T2 | Europe, thiabendazole at 3.0 kg over the rows at sowing in 1982 and in spring 1983 and 1984 |

NOTE: Treatments in 1984 were applied in 7500 l by weeder bar.

Basal applications: Manures: (0:24:24) at 730 kg. Weedkiller: Propyzamide at 0.70 kg in 220 l.

Cultivations, etc.:-

Both sites: Weedkiller applied: 18 Jan, 1984. PK applied: 15 Mar.

Healthy site: Cut: 12 June. Aldicarb and thiabendazole treatments applied: 21 June. Cut: 17 July. Aldicarb applied: 23 July. Cut: 3 Sept.

Infested site: Cut: 14 June. Aldicarb applied: 21 June. Thiabendazole applied: 28 June. Cut: 6 Aug. Aldicarb applied: 10 Aug. Cut: 26 Sept.

84/R/CS/279

NOTE: Assessments of stems infected with stem nematode were made on both sites.

LONG HOOS V 5 (HEALTHY SITE)

1ST CUT (12/6/84) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| TREATMNT | |
|----------|------|
| V 0 | 6.45 |
| V A1 | 6.63 |
| E 0 | 7.39 |
| E A1 | 7.40 |
| E A2 | 7.24 |
| E A1 A1 | 7.31 |
| E A1 T1 | 6.73 |
| E A2 T2 | 6.63 |
| E C1 | 6.73 |
| E C2 | 7.30 |
| E C1 T1 | 6.54 |
| E C2 T2 | 7.24 |
| E T1 T1 | 6.61 |
| E T2 T2 | 6.86 |
| MEAN | 6.93 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | TREATMNT |
|-------|----------|
| ----- | ----- |
| SED | 0.575 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|--------------|------|-------|------|
| BLOCK.WP | 26 | 0.705 | 10.2 |
| 1ST MEAN DM% | 15.1 | | |

84/R/CS/279 LONG HOOS V 5 (HEALTHY SITE)

2ND CUT (17/7/84) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| TREATMNT | |
|----------|------|
| V 0 | 4.27 |
| V A1 | 4.23 |
| E 0 | 5.17 |
| E A1 | 5.05 |
| E A2 | 5.35 |
| E A1 A1 | 4.73 |
| E A1 T1 | 4.95 |
| E A2 T2 | 4.75 |
| E C1 | 4.91 |
| E C2 | 4.84 |
| E C1 T1 | 5.32 |
| E C2 T2 | 4.79 |
| E T1 T1 | 5.05 |
| E T2 T2 | 4.52 |
| MEAN | 4.85 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | TREATMNT |
|-------|----------|
| ----- | ----- |
| SED | 0.247 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|--------------|----|-------|-----|
| BLOCK.WP | 26 | 0.303 | 6.2 |
| 2ND MEAN DM% | | 17.9 | |

84/R/CS/279 LONG HOOS V 5 (HEALTHY SITE)
 3RD CUT (3/9/84) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| TREATMNT | |
|----------|------|
| V 0 | 3.86 |
| V A1 | 3.72 |
| E 0 | 4.48 |
| E A1 | 4.36 |
| E A2 | 4.34 |
| E A1 A1 | 4.25 |
| E A1 T1 | 4.58 |
| E A2 T2 | 4.36 |
| E C1 | 4.63 |
| E C2 | 4.38 |
| E C1 T1 | 4.92 |
| E C2 T2 | 4.33 |
| E T1 T1 | 4.71 |
| E T2 T2 | 4.35 |
| MEAN | 4.38 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | TREATMNT |
|-------|----------|
| ----- | ----- |
| SED | 0.264 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|------------------|------|-------|-----|
| BLOCK.WP | 26 | 0.324 | 7.4 |
| 3RD CUT MEAN DM% | 22.8 | | |

84/R/CS/279 LONG HOOS V 5 (HEALTHY SITE)

TOTAL OF 3 CUTS DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| TREATMNT | |
|----------|-------|
| V 0 | 14.58 |
| V A1 | 14.58 |
| E 0 | 17.04 |
| E A1 | 16.81 |
| E A2 | 16.93 |
| E A1 A1 | 16.29 |
| E A1 T1 | 16.25 |
| E A2 T2 | 15.73 |
| E C1 | 16.27 |
| E C2 | 16.52 |
| E C1 T1 | 16.78 |
| E C2 T2 | 16.35 |
| E T1 T1 | 16.37 |
| E T2 T2 | 15.73 |
| MEAN | 16.16 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | TREATMNT |
|-------|----------|
| ----- | ----- |
| SED | 0.622 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|----------|----|-------|-----|
| BLOCK.WP | 26 | 0.762 | 4.7 |

TOTAL OF 2 CUTS MEAN DM% 18.6

PLOT AREA HARVESTED 0.00045

84/R/CS/279 LONG HOOS IV 2 (INFESTED SITE)

1ST CUT (14/6/84) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| TREATMNT | |
|----------|------|
| V 0 | 5.30 |
| V A1 | 5.10 |
| E 0 | 5.46 |
| E A1 | 4.13 |
| E A2 | 4.29 |
| E A1 A1 | 5.02 |
| E A1 T1 | 3.55 |
| E A2 T2 | 4.04 |
| E C1 | 4.91 |
| E C2 | 5.51 |
| E C1 T1 | 4.72 |
| E C2 T2 | 4.40 |
| E T1 T1 | 4.88 |
| E T2 T2 | 4.74 |
| MEAN | 4.72 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | TREATMNT |
|-------|----------|
| ----- | ----- |
| SED | 0.473 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|--------------|------|-------|------|
| BLOCK.WP | 26 | 0.580 | 12.3 |
| 1ST MEAN DM% | 15.8 | | |

84/R/CS/279 LONG HOOS IV 2 (INFESTED SITE)

2ND CUT (6/8/84) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| TREATMNT | |
|----------|------|
| V 0 | 7.39 |
| V A1 | 5.59 |
| E 0 | 6.41 |
| E A1 | 4.64 |
| E A2 | 5.95 |
| E A1 A1 | 5.97 |
| E A1 T1 | 5.50 |
| E A2 T2 | 5.27 |
| E C1 | 6.33 |
| E C2 | 6.29 |
| E C1 T1 | 5.16 |
| E C2 T2 | 5.91 |
| E T1 T1 | 5.64 |
| E T2 T2 | 5.24 |
| MEAN | 5.81 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | TREATMNT |
|-------|----------|
| ----- | ----- |
| SED | 0.678 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|--------------|------|-------|------|
| BLOCK.WP | 26 | 0.830 | 14.3 |
| 2ND MEAN DM% | 24.7 | | |

84/R/CS/279 LONG HOOS IV 2 (INFESTED SITE)
3RD CUT (26/9/84) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| TREATMNT | |
|----------|------|
| V 0 | 2.98 |
| V A1 | 2.01 |
| E 0 | 2.47 |
| E A1 | 1.95 |
| E A2 | 2.30 |
| E A1 A1 | 2.08 |
| E A1 T1 | 1.99 |
| E A2 T2 | 2.00 |
| E C1 | 2.46 |
| E C2 | 2.34 |
| E C1 T1 | 2.48 |
| E C2 T2 | 1.91 |
| E T1 T1 | 2.34 |
| E T2 T2 | 2.03 |
| MEAN | 2.24 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | TREATMNT |
|-------|----------|
| ----- | ----- |
| SED | 0.326 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|----------|----|-------|------|
| BLOCK.WP | 26 | 0.399 | 17.8 |

3RD CUT MEAN DM% 22.2

84/R/CS/279 LONG HOOS IV 2 (INFESTED SITE)

TOTAL OF 3 CUTS DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| TREATMNT | |
|----------|-------|
| V 0 | 15.68 |
| V A1 | 12.69 |
| E 0 | 14.34 |
| E A1 | 10.73 |
| E A2 | 12.54 |
| E A1 A1 | 13.06 |
| E A1 T1 | 11.04 |
| E A2 T2 | 11.31 |
| E C1 | 13.70 |
| E C2 | 14.14 |
| E C1 T1 | 12.36 |
| E C2 T2 | 12.22 |
| E T1 T1 | 12.86 |
| E T2 T2 | 12.01 |
| MEAN | 12.76 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | TREATMNT |
|-------|----------|
| ----- | ----- |
| SED | 1.294 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|----------|----|-------|------|
| BLOCK.WP | 26 | 1.585 | 12.4 |

TOTAL OF 2 CUTS MEAN DM% 20.9

PLOT AREA HARVESTED 0.00045