

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1984

[Full Table of Content](#)



## 84/R/WW/6 Seed Dressings and Take-all - W. Wheat

### Rothamsted Research

Rothamsted Research (1985) *84/R/WW/6 Seed Dressings and Take-all - W. Wheat* ; Yields Of The Field Experiments 1984, pp 232 - 233 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/WW/6

WINTER WHEAT

SEED DRESSINGS AND TAKE-ALL

Object: To study the effects of seed treatments on the incidence of take-all and on the yield of early-sown winter wheat - Gt. Knott I.

Sponsor: G.L. Bateman.

Design: 4 randomised blocks of 6 plots.

Whole plot dimensions: 3.0 x 12.0.

Treatments: All combinations of:-

1. SOWDATE            Dates of sowing:  
    8 SEPT            8 September, 1983  
    7 OCT             7 October
2. SEED DR            Seed dressings:  
    NONE             None  
    TRIADIME         Triadimenol at 0.5 g/kg of seed plus fuberidazole at  
                            0.06 g/kg of seed  
    UKO 82F          'UKO 82 f' at 2.75 ml/kg of seed

Basal applications: Manures: Muriate of potash at 420 kg. 'Nitro-Chalk' on three occasions, at 130 kg on the first, at 150 kg on the second and at 460 kg on the third. Weedkillers: Isoproturon at 2.5 kg with bromoxynil and ioxynil (as 'Deloxil' at 2.0 l) in 250 l. Dicamba with mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l applied with the prochloraz and carbendazim. Fungicides: Prochloraz at 0.4 kg with carbendazim at 0.15 kg. Maneb at 1.6 kg with tridemorph at 0.38 kg and carbendazim at 0.15 kg in 250 l. Insecticide: Pirimicarb at 0.14 kg in 250 l.

Seed: Aquila, sown at 170 kg.

Cultivations, etc.:- Heavy spring-tine cultivated: 3 Sept, 1983. Muriate of potash and first N applied: 7 Sept. All plots heavy spring-tine cultivated twice, early-sown plots rotary harrowed and seed sown: 8 Sept. Late-sown plots rotary harrowed and seed sown: 7 Oct. Isoproturon with 'Deloxil' applied: 1 Dec. Second N applied: 16 Feb, 1984. Third N applied: 11 Apr. 'Herrisol' with prochloraz and carbendazim applied: 17 Apr. Maneb, carbendazim and tridemorph applied: 13 June. Insecticide applied: 27 June. Combine harvested: 22 Aug. Previous crops: W. wheat 1982 and 1983.

NOTE: Take-all and mildew assessments were made twice during the autumn. Take-all and foot rots were assessed in March, May and June.

84/R/WW/6

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

SEED DR SOWDATE	NONE	TRIADIME	UKO 82F	MEAN
8 SEPT	5.72	8.44	6.93	7.03
7 OCT	7.43	7.75	7.89	7.69
MEAN	6.57	8.09	7.41	7.36

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	SOWDATE	SEED DR	SOWDATE SEED DR
SED	0.378	0.463	0.655

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	15	0.926	12.6

GRAIN MEAN DM% 86.2

PLOT AREA HARVESTED 0.00329