

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/WW/15 Chloride and Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (1985) *84/R/WW/15 Chloride and Take-all - W. Wheat* ; Yields Of The Field Experiments 1984, pp 240 - 241 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/WW/15

WINTER WHEAT

CHLORIDE AND TAKE-ALL

Object: To study the effects of different spring nitrogen top dressings, that include chloride and ammonium ions, on the incidence of take-all and on the yield of winter wheat - Gt. Knott I.

Sponsors: R.J. Gutteridge, G.L. Bateman.

Design: 4 randomised blocks of 4 plots.

Whole plot dimensions: 3.0 x 12.0.

Treatments:

SPRING N Spring nitrogen, 40 kg N on 9 March, 1984; 160 kg N on 16 April:

AMM CHL	Ammonium chloride
AMM NIT	Ammonium nitrate as 'Nitro-Chalk'
AMM SUL	Ammonium sulphate
UREA	Urea

Basal applications: Manures: Muriate of potash at 410 kg. 'Nitro-Chalk' at 130 kg. Weedkillers: Isoproturon at 2.5 kg with bromoxynil and ioxynil (as 'Deloxil' at 1.0 l) in 250 l. Dicamba, mecoprop and MCPA (as 'Herrisol' at 5.0 l) in 250 l applied with the prochloraz and carbendazim. Fungicides: Prochloraz at 0.40 kg and carbendazim at 0.15 kg. Maneb at 1.6 kg with carbendazim at 0.15 kg and tridemorph at 0.38 kg in 250 l. Insecticide: Pirimicarb at 0.14 kg in 250 l.

Seed: Longbow, sown at 170 kg.

Cultivations, etc.: - Discd: 3 Sept, 1983. Basal N and K applied: 7 Sept. Rotary harrowed, seed sown: 11 Oct. Isoproturon with 'Deloxil' applied: 30 Nov. 'Herrisol' with prochloraz and carbendazim applied: 17 Apr, 1984. Maneb with carbendazim and tridemorph applied: 13 June. Insecticide applied: 27 June. Combine harvested: 22 Aug. Previous crops: W. wheat 1982 and 1983.

NOTE: Take-all assessments were made monthly from early March to late June and foot rot assessments were made in June.

84/R/WW/15

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

SPRING N	AMM CHL	AMM NIT	AMM SUL	UREA	MEAN
	7.92	8.59	8.25	8.33	8.27

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT
-----	-----
SED	0.453

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	11	0.641	7.8

GRAIN MEAN DM% 88.0

PLOT AREA HARVESTED 0.00280