

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/B/2 Electrostatic Spraying and Foliar Diseases - W. Barley

Rothamsted Research

Rothamsted Research (1985) *84/R/B/2 Electrostatic Spraying and Foliar Diseases - W. Barley* ; Yields Of The Field Experiments 1984, pp 260 - 261 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/B/2

WINTER BARLEY

ELECTROSTATIC SPRAYING AND FOLIAR DISEASES

Object: To study the penetration of sprays and control of foliar diseases with a range of electrostatic sprayers - Black Horse II.

Sponsors: D.C. Griffiths, G.R. Cayley, B.J. Pye, P. Etheridge, G.C. Scott, F.T. Phillips.

Design: 4 randomised blocks of 8 plots.

Whole plot dimensions: 3.0 x 15.0.

Treatments:

SPRAYER	Sprayers applying propiconazole:
NONE	None
CNVNTL 2	Conventional hydraulic sprayer, at 125 g in 200 l
CNVNTL 1	Conventional hydraulic sprayer, at 62.5 g in 200 l
EL APE	'APE' electrostatic sprayer, at 62.5 g in 6 l (duplicated)
EL JUMBO	'Jumbo' electrostatic sprayer, at 62.5 g in 10 l (duplicated)
EL MICRO	'Micronex' electrostatic sprayer, at 62.5 g in 13 l

- NOTES: (1) Propiconazole was applied on 3 November, 1983 and 14 March, 1984 by all sprayers except the 'Micronex' which was on 14 March only.
- (2) The 'APE' electrostatic sprayer had four spinning-disc nozzles mounted on a hand-held boom, the 'Jumbo' had spinning-cone nozzles. Both are charged at 30 kv.
- (3) The 'Micronex' is a commercial prototype, electrostatically-charged spinning-disc sprayer.
- (4) Chopped straw infected with *Rhynchosporium* was spread evenly over the whole of the experimental area on 9 September, 1983.

Basal applications: Manures: (5:14:30) at 340 kg. 'Nitro-Chalk' on two occasions, at 190 kg on the first and at 440 kg on the second.

Weedkillers: Methabenzthiazuron at 2.4 kg in 250 l. 3, 6-dichloropicolinic acid at 0.07 kg and bromoxynil at 0.34 kg with mecoprop (as 'CMPP' at 4.2 l) in 200 l. Desiccant: Diquat at 0.70 kg ion with 'Agral', a wetting agent, at 0.2 l, in 200 l.

Seed: Maris Otter, sown at 160 kg.

Cultivations, etc.: - Ploughed: 4 Aug, 1983. NPK applied: 23 Aug. Spring-tine cultivated: 7 Sept. Straw applied, rotary harrowed, seed sown: 9 Sept. Methabenzthiazuron applied: 13 Sept. First N applied: 9 Mar, 1984. Second N applied: 4 Apr. 3, 6-dichloropicolinic acid, bromoxynil and mecoprop applied: 13 Apr. Desiccant applied: 23 July. Combine harvested: 26 July. Previous crops: W. barley 1982 and 1983.

NOTE: Plant samples were taken immediately after spraying to assess weedkiller deposits. Mildew was assessed in November and *Rhynchosporium* in April.

84/R/B/2

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

SPRAYER	NONE	CNVNTL 2	CNVNTL 1	EL APE	EL JUMBO	EL MICRO	MEAN
	5.96	6.65	6.64	6.64	6.53	6.27	6.48

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	SPRAYER
-----	-----
SED	0.238 MIN REP
	0.206 MAX-MIN
	0.168 MAX REP

	SPRAYER
MAX REP	EL APE V EL JUMBO
MAX-MIN	EL APE OR EL JUMBO V ANY OF REMAINDER
MIN REP	ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	23	0.336	5.2

GRAIN MEAN DM% 78.6

PLOT AREA HARVESTED 0.00306