

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/W/B/2 Sowing Dates and Insects - S. Barley

Rothamsted Research

Rothamsted Research (1985) *84/W/B/2 Sowing Dates and Insects - S. Barley* ; Yields Of The Field Experiments 1984, pp 262 - 263 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/W/B/2

SPRING BARLEY

SOWING DATES AND INSECTS

Object: To study the effects of omethoate on insect pests and on yields of s. barley sown on two dates - Woburn White Horse.

Sponsor: G.C. Scott.

Design: 4 randomised blocks of 8 plots.

Whole plot dimensions: 8.0 x 12.0.

Treatments: All combinations of:-

- | | |
|-------------|----------------------------|
| 1. SOW DATE | Dates of sowing: |
| 9 MAR | 9 Mar, 1984 |
| 16 APR | 16 Apr |
| 2. INSEARLY | Insecticide applied early: |
| NONE | None |
| OMETHO E | Omethoate on 31 May |
| 3. INS LATE | Insecticide applied late: |
| NONE | None |
| OMETHO L | Omethoate on 29 June |

NOTE: Omethoate was applied at 0.63 l in 250 l on both occasions.

Basal applications: Manures: Magnesian limestone at 7.5 t, FYM at 50 t, N at 110 kg as 'Nitro-Chalk'. Weedkiller: Mecoprop at 2.1 kg in 250 l. Fungicide: Tridemorph at 0.52 kg in 250 l.

Seed: Triumph, dressed with triadimenol plus fuberidazole, sown at 160 kg.

Cultivations, etc.:- Magnesian limestone applied: 30 Sept, 1983. FYM applied: 21-23 Nov. Ploughed: 25 Nov. N applied: 8 Mar, 1984. Spring-tine cultivated all plots, spring-tine cultivated with crumbler attached, seed sown for SOW DATE 9 MAR: 9 Mar. Spring-tine cultivated with crumbler attached, seed sown for SOW DATE 16 APR: 16 Apr. Weedkiller applied: 15 May. Fungicide applied: 31 May. Combine harvested: 15 Aug. Previous crops: Potatoes 1982, w. wheat 1983.

NOTES: (1) Aphids, thrips and stem borers were counted on several occasions during the growing season.
(2) Barley yellow dwarf virus infection was assessed.

84/W/B/2

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

INSEARLY SOWDATE	NONE	OMETHO E	MEAN
9 MAR	7.13	6.81	6.97
16 APR	5.55	5.99	5.77
MEAN	6.34	6.40	6.37

INS LATE SOWDATE	NONE	OMETHO L	MEAN
9 MAR	6.90	7.04	6.97
16 APR	5.58	5.97	5.77
MEAN	6.24	6.51	6.37

INS LATE INSEARLY	NONE	OMETHO L	MEAN
NONE	6.20	6.49	6.34
OMETHO E	6.28	6.52	6.40
MEAN	6.24	6.51	6.37

INSEARLY INS LATE SOWDATE	NONE	OMETHO L	OMETHO E NONE	OMETHO L
9 MAR	7.19	7.07	6.61	7.01
16 APR	5.20	5.90	5.95	6.04

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	SOWDATE	INSEARLY	INS LATE	SOWDATE INSEARLY
SED	0.265	0.265	0.265	0.375

TABLE	SOWDATE INS LATE	INSEARLY INS LATE	SOWDATE INSEARLY INS LATE
SED	0.375	0.375	0.531

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	21	0.751	11.8

GRAIN MEAN DM% 87.1

PLOT AREA HARVESTED 0.00275