

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/B/14 Mildew Control by Electrostatic Sprayers - S. Barley

Rothamsted Research

Rothamsted Research (1985) *84/R/B/14 Mildew Control by Electrostatic Sprayers - S. Barley* ; Yields Of The Field Experiments 1984, pp 278 - 279 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/B/14

SPRING BARLEY

MILDEW CONTROL BY ELECTROSTATIC SPRAYERS

Object: To compare the effects of a range of electrostatic sprayers on mildew (*Erysiphe graminis*) control and on the yield of spring barley - Webbs.

Sponsors: D.C. Griffiths, G.R. Cayley, P. Etheridge, R.E. Goodchild, B.J. Pye, G.C. Scott.

Design: 4 randomised blocks of 14 plots.

Whole plot dimensions: 3.0 x 15.0.

Treatments:

SPRAYER	Sprayers and rates of application of propiconazole (kg):
CNVNTL 1	Conventional, 0.125 kg
CNVNTL 2	Conventional, 0.250 kg
	Electrostatic sprayers, applying at 0.125 kg:
E NO 1	NIAE, uncharged hydraulic sprayer in 60 l
E NC 1	NIAE, charged hydraulic sprayer in 60 l
E JC 1	'Jumbo', charged in 10.1 l
E AC 1	'APE' charged in 5.6 l
E MMC 1	'Micromax' charged in 9.0 l
E MNC 1	'Micronex' charged in 9.0 l
E JAAC 1	'Jumbo', air assisted, nozzles mounted at 20° to vertical, charged, in 10.1 l
E JAVC 1	'Jumbo', air assisted, nozzles mounted vertically, charged, in 10.1 l
E AAAC 1	'APE', air assisted, nozzles mounted at 20° to vertical, charged, in 5.6 l
E AAVC 1	'APE', air assisted, nozzles mounted vertically, charged, in 5.6 l
NONE	None (duplicated)

- NOTES: (1) Sides and ends of plots, 3m and 5m respectively, were separated by *Atem s.* barley, seed dressed with triadimenol plus fuberidazole.
- (2) The 'Jumbo' has electrostatically charged spinning-cone nozzles.
- (3) The 'APE' has electrostatically charged spinning-disc nozzles.
- (4) The 'Micromax' has inductively-charged spinning-cone nozzles.
- (5) The 'Micronex' is a commercial prototype, electrostatically-charged spinning-disc sprayer.
- (6) Spray treatments were applied in the period 11 to 12 June, 1984.

Basal applications: Manures: FYM at 25 t. 'Nitro-Chalk' at 500 kg.
Weedkillers: 3, 6-dichloropicolinic acid at 0.05 kg and bromoxynil at 0.24 kg with mecoprop (as 'CMPP' at 3.0 l) in 250 l.

Seed: Georgie, sown at 160 kg.

84/R/B/14

Cultivations, etc.: - Heavy spring-tine cultivated: 10 Oct, 1983. FYM applied: 11 Nov. Ploughed: 28 Nov. N applied, spring-tine cultivated: 8 Mar, 1984. Rotary harrowed, seed sown: 10 Mar. Weedkillers applied: 15 May. Combine harvested: 20 Aug. Previous crops: S. barley 1982 and 1983.

NOTE: Plant samples were taken immediately after spraying to assess weedkiller deposits. Mildew assessments were made in June.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

SPRAYER	
CNVNTL 1	6.30
CNVNTL 2	6.19
E NO 1	5.96
E NC 1	6.20
E JC 1	5.65
E AC 1	5.74
E MMC 1	5.72
E MNC 1	5.83
E JAAC 1	6.19
E JAVC 1	6.12
E AAAC 1	5.92
E AAVC 1	6.19
NONE	5.42
MEAN	5.92

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	SPRAYER
-----	-----
SED	0.185 MIN REP
	0.160 MAX-MIN

SPRAY
 MAX-MIN NONE V ANY OF REMAINDER
 MIN REP ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	40	0.261	4.4
GRAIN MEAN DM%	87.0		
PLOT AREA HARVESTED	0.00306		