

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/BE/1 Effects of Pests and Pathogens - W. Beans

Rothamsted Research

Rothamsted Research (1985) *84/R/BE/1 Effects of Pests and Pathogens - W. Beans* ; Yields Of The Field Experiments 1984, pp 280 - 281 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/BE/1

WINTER BEANS

EFFECTS OF PESTS AND PATHOGENS

Object: To assess the effects of three amounts of pest and disease control on w. beans - Highfield IV.

Sponsors: J. McEwen, R. Bardner, A.J. Cockbain, D.C. Griffiths, D.H. Lapwood, R.M. Webb, D.P. Yeoman.

Design: 6 randomised blocks of 3 plots.

Whole plot dimensions: 5.33 x 15.0.

Treatments:

PATHCONT	Pest and pathogen control (in addition to basals):
STANDARD	None
ENHANCED	Seed dressed with carbendazim and thiram (1.1 g of each per kg of seed)
FULL	Phorate at 1.7 kg as granules to foliage on 12 Apr, 1984 Seed dressed with carbendazim and thiram Aldicarb at 10 kg on 29 Sept, 1983 Fosetyl-Al at 1.6 kg on 14 Mar, 1984 Benomyl at 0.56 kg and chlorothalonil at 0.98 kg in 340 l on 4 Apr Carbofuran at 1.7 kg on 12 Apr Propiconazole at 0.12 kg on 4 July

NOTES: (1) Treatment sprays were applied in 220 l except where stated.
(2) Sides of plots were separated by strips of w. beans 5.33 m wide plus 0.66 m fallow paths, ends of plots were separated by strips of w. beans 9.2 m wide plus 0.9 m fallow paths. The separating crops received basal applications as for the plots.

Basal applications: Weedkillers: Paraquat at 0.42 kg ion in 250 l. Propyzamide at 0.85 kg with simazine at 1.2 l in 250 l. Fungicides: Benomyl at 0.50 kg with chlorothalonil at 1.0 kg and 'Agral', a wetting agent at 0.08 l, in 250 l on the first two occasions and in 200 l on the third occasion.

Seed: Banner, sown at 220 kg.

Cultivations, etc.: Heavy spring-tine cultivated: 1 Sept, 1983. Chisel ploughed twice: 2 Sept. Paraquat applied: 20 Sept. Heavy spring-tine cultivated: 24 Sept. Aldicarb treatment applied, rotary harrowed, seed sown: 29 Sept. Propyzamide and simazine applied: 30 Sept. Basal fungicides applied: 24 May, 5 June, 26 June, 1984. Combine harvested: 30 Aug. Previous crops: W. wheat 1982, s. barley 1983.

NOTE: Plant counts were made after establishment and components of yield were measured at maturity. Migratory nematodes, root and foliar fungi, aphids and weevils were counted at intervals during the season. Total above-ground dry matter and N content were measured in July. N content of grain was measured.

84/R/BE/1

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PATHCONT	STANDARD	ENHANCED	FULL	MEAN
	4.15	4.34	4.26	4.25

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PATHCONT
-------	----------

SED	0.230
-----	-------

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.398	9.4

GRAIN MEAN DM% 85.9

PLOT AREA HARVESTED 0.00320