

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/BE/16 Control of Rust - S. Beans

Rothamsted Research

Rothamsted Research (1985) *84/R/BE/16 Control of Rust - S. Beans ; Yields Of The Field Experiments 1984*, pp 303 - 306 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/BE/16

SPRING BEANS

CONTROL OF RUST

Object: To study the effects of fungicides on the control of rust (*Uromyces viciae-fabae*) and on the yield of unirrigated and irrigated s. beans - Long Hoos IV 3.

Sponsors: D.H. Lapwood, J. McEwen, D.P. Yeoman.

Design: 2 randomised blocks of 2 plots split into 12.

Whole plot dimensions: 2.03 x 2.13.

Treatments: All combinations of:-

Whole plots

1. IRRIGATN	Irrigation:
0	None
I	Irrigated (86 mm)

Sub plots

2. C S FUNG	Fungicide to control chocolate spot but not rust:
NONE	None
BENOMYL	Benomyl at 0.56 kg in 220 l on 6 July, 14 Aug, 1984
3. RUSTFUNG	Fungicides to control rust:
MAN+MANC	Maneb at 0.8 kg + mancozeb at 0.8 kg in 340 l
PROPICON	Propiconazole at 0.12 kg in 340 l
4. RFNGTIME	Times of applying fungicides to control rust:
ONCE	Once on 1 Aug
TWICE	Twice, on 1 Aug and 8 Aug

plus two extra sub plot treatments:

EXTRA

NONE	No fungicides (duplicated)
BENOMYL	Benomyl at 0.56 kg in 220 l on 6 July, 14 Aug (duplicated)

NOTE: After two post-flowering applications totalling 38 mm irrigation was applied subsequently at 8 mm, on two occasions per week, unless 4 mm or more of rain had fallen since the last application.

84/R/BE/16

Date	mm water
13 July	13
20 July	25
24 July	8
27 July	8
31 July	8
3 Aug	8
21 Aug	8
24 Aug	8
—	
Total	86 mm

Basal applications: Manures: Chalk at 2.9 t. Weedkillers: Trietazine at 0.95 kg with simazine at 0.14 kg in 220 l. Insecticides: Permethrin at 0.15 kg in 220 l; cypermethrin at 0.025 kg in 220 l on two occasions; pirimicarb at 0.14 kg in 220 l on two occasions.

Seed: Minden, sown at 200 kg.

Cultivations, etc.:— Chalk applied: 25 Aug, 1983. Ploughed: 17 Nov. Spring-tine cultivated, seed sown: 19 Mar, 1984. Weedkillers applied: 2 Apr. Permethrin applied: 1 May. Cypermethrin applied: 16 May, 6 June. Pirimicarb applied: 4 June, 6 July. Harvested by hand: 23 Aug (unirrigated plots), 30 Aug (irrigated plots). Previous crops: Potatoes 1982, s. wheat 1983.

NOTE: Plant counts were made after establishment. The incidence of chocolate spot and rust were assessed from early July until maturity. Components of yield were measured at maturity.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

C S FUNG IRRIGATN	NONE	BENOMYL	MEAN
0	4.78	4.47	4.63
I	5.37	5.27	5.32
MEAN	5.08	4.87	4.97
RUSTFUNG IRRIGATN	MAN+MANC	PROPICON	MEAN
0	4.77	4.48	4.63
I	5.43	5.21	5.32
MEAN	5.10	4.85	4.97

84/R/BE/16

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

RUSTFUNG C S FUNG	MAN+MANC	PROPICON	MEAN
NONE	5.35	4.80	5.08
BENOMYL	4.85	4.89	4.87
MEAN	5.10	4.85	4.97

RFNGTIME IRRIGATN	ONCE	TWICE	MEAN
0	4.86	4.40	4.63
I	5.31	5.32	5.32
MEAN	5.09	4.86	4.97

RFNGTIME C S FUNG	ONCE	TWICE	MEAN
NONE	5.22	4.93	5.08
BENOMYL	4.95	4.78	4.87
MEAN	5.09	4.86	4.97

RFNGTIME RUSTFUNG MAN+MANC PROPICON	ONCE	TWICE	MEAN
MAN+MANC	5.26	4.94	5.10
PROPICON	4.91	4.78	4.85
MEAN	5.09	4.86	4.97

C S FUNG RUSTFUNG IRRIGATN	NONE MAN+MANC	PROPICON	BENOMYL MAN+MANC	PROPICON
0	5.09	4.48	4.45	4.49
I	5.61	5.13	5.24	5.29

C S FUNG RFNGTIME IRRIGATN	NONE ONCE	TWICE	BENOMYL ONCE	TWICE
0	5.04	4.53	4.68	4.26
I	5.41	5.33	5.22	5.31

RUSTFUNG RFNGTIME IRRIGATN	MAN+MANC ONCE	TWICE	PROPICON ONCE	TWICE
0	5.06	4.49	4.66	4.31
I	5.46	5.39	5.17	5.25

RUSTFUNG RFNGTIME C S FUNG	MAN+MANC ONCE	TWICE	PROPICON ONCE	TWICE
NONE	5.42	5.28	5.02	4.59
BENOMYL	5.09	4.60	4.81	4.97

84/R/BE/16

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

IRRIGATN	C S FUNG	RUSTFUNG	MAN+MANC	PROPICON	
		RFNGTIME	ONCE	TWICE	ONCE
0	NONE	5.29	4.89	4.78	4.17
	BENOMYL	4.82	4.08	4.54	4.44
I	NONE	5.56	5.66	5.25	5.00
	BENOMYL	5.36	5.12	5.08	5.50
EXTRA IRRIGATN	NONE	BENOMYL	MEAN		
0	4.41	4.45	4.43		
I	5.24	5.41	5.32		
MEAN	4.82	4.93	4.87		

GRAND MEAN 4.94

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	C S FUNG	RUSTFUNG	RFNGTIME
SED	0.140	0.099	0.099	0.099
TABLE	IRRIGATN* C S FUNG	IRRIGATN* RUSTFUNG	C S FUNG RUSTFUNG	IRRIGATN* RFNGTIME
SED	0.140	0.140	0.140	0.140
TABLE	C S FUNG RFNGTIME	RUSTFUNG RFNGTIME	IRRIGATN* C S FUNG RUSTFUNG	IRRIGATN* C S FUNG RFNGTIME
SED	0.140	0.140	0.198	0.198
TABLE	IRRIGATN* RUSTFUNG RFNGTIME	C S FUNG RUSTFUNG RFNGTIME	IRRIGATN* C S FUNG RUSTFUNG RFNGTIME	IRRIGATN* EXTRA
SED	0.198	0.198	0.281	0.198

* WITHIN THE SAME LEVEL OF IRRIGATN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	26	0.281	5.7
GRAIN MEAN DM%	89.2		
PLOT AREA HARVESTED	0.00015		