

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1984

[Full Table of Content](#)



## 84/R/FE/1 N, *Rhizobium* and Pest Control - Fenugreek

### Rothamsted Research

Rothamsted Research (1985) *84/R/FE/1 N, Rhizobium and Pest Control - Fenugreek* ; Yields Of The Field Experiments 1984, pp 311 - 312 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/FE/1

FENUGREEK

N, RHIZOBIUM AND PEST CONTROL

Object: To study the effects of inoculation with Rhizobium, application of insecticide and times of applying nitrogen fertilizer on nodulation and yield of fenugreek (*Trigonella foenum - graecum*) - Long Hoos III 7.

Sponsor: D.P. Yeoman.

Design: 2 randomised blocks of 14 plots.

Whole plot dimensions: 2.64 x 8.00.

Treatments: All combinations of:-

1. INOCULUM            Inoculum applied to the seed:  
    NONE                None  
    RHIZOBUM           Rhizobium meliloti, strain 2012, as a peat culture
2. N                    Nitrogen fertilizer (kg N) and times of application:  
    0                    None  
    150 S                150 to the seedbed, on 14 May, 1984  
    150 F                150 at flowering, on 16 July
3. INSECTICIDE        Insecticide:  
    NONE                None  
    PERMETH            Permethrin foliar spray at 0.053 kg in 220 l on 15 June and 6 July

plus two extra treatments, each given 30 kg N to the seedbed and permethrin as above:

EXTRA

- |          |                             |
|----------|-----------------------------|
| O N30S P | No inoculum                 |
| R N30S P | Rhizobium inoculum as above |

Basal applications: Manures: Chalk at 2.9 t. Weedkillers: Glyphosate at 0.72 kg in 200 l. MCPB at 2.24 kg in 220 l. Trifluralin at 0.81 kg in 220 l. Desiccant: Diquat at 0.84 kg ion in 220 l on two occasions.

Seed: Barbara, sown at 29 kg.

Cultivations, etc.:- Chalk applied: 26 Aug, 1983. Glyphosate applied: 16 Dec. Ploughed: 30 Jan, 1984. Spring-tine cultivated, trifluralin applied, power harrowed, seed sown: 14 May. MCPB applied: 5 July. Hand weeded: 24 July. Desiccant applied: 7 Sept, 12 Sept. Combine harvested: 27 Sept. Previous crops: S. wheat 1982, mixed cereals 1983.

NOTE: Plant counts were made after establishment. N content of grain was measured.

84/R/FE/1

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	N	0	150 S	150 F	MEAN
INOCULUM					
NONE		0.05	0.11	0.07	0.08
RHIZOBUM		0.09	0.11	0.07	0.09
MEAN		0.07	0.11	0.07	0.08

	INSCTCDE	NONE	PERMETH	MEAN
INOCULUM				
NONE		0.06	0.09	0.08
RHIZOBUM		0.09	0.09	0.09
MEAN		0.08	0.09	0.08

	INSCTCDE	NONE	PERMETH	MEAN
N				
0		0.06	0.08	0.07
150 S		0.08	0.14	0.11
150 F		0.08	0.05	0.07
MEAN		0.08	0.09	0.08

	N	0	150 S	150 F
INSCTCDE				
NONE		0.05	0.05	0.09
RHIZOBUM		0.08	0.11	0.07
NONE			0.05	0.17
RHIZOBUM			0.12	0.11
NONE				0.09
RHIZOBUM				0.05
NONE				0.07
RHIZOBUM				0.06

EXTRA	0	N30S	P	R	N30S	P	MEAN
			0.09			0.10	0.09

GRAND MEAN 0.09

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	EXTRA	INOCULUM	N	INSCTCDE
SED	0.043	0.018	0.022	0.018

TABLE	INOCULUM	INOCULUM	N	INOCULUM
	N	INSCTCDE	INSCTCDE	N
	INSCTCDE & EXTRA			
SED	0.031	0.025	0.031	0.043

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	13	0.043	50.5

GRAIN MEAN DM% 55.9  
 PLOT AREA HARVESTED 0.00154