

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/RA/2 Urea and Inhibitors - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1985) *84/R/RA/2 Urea and Inhibitors - W. Oilseed Rape* ; Yields Of The Field Experiments 1984, pp 313 - 315 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/RA/2

WINTER OILSEED RAPE

UREA AND INHIBITORS

Object: To study the effects of adding nitrification inhibitors to prilled urea, applied to the seedbed and in spring on the yield and nitrogen uptake of w. oilseed rape - Whittlocks.

Sponsors: G.A. Rodgers, A. Penny, M.V. Hewitt.

Design: 2 randomised blocks of 17 plots.

Whole plot dimensions: 4.0 x 20.0.

Treatments: All combinations of:-

1. N INHIB Forms of nitrogen and nitrification inhibitor used for seedbed and spring nitrogen applications:

AN 0	Ammonium nitrate (as 'Nitro-Chalk'), no inhibitor
PU 0	Prilled urea, no inhibitor
PU DIC	Prilled urea and dicyandiamide
PU HYD	Prilled urea and hydroquinone
2. SEEDBD N Nitrogen rates (kg N) to seedbed (on 22 August, 1983):

0	
50	
3. SPRING N Nitrogen rates (kg N) and times in spring:

75E+75L	75 on 10 Feb and 75 on 21 Mar, 1984.
150M	150 on 5 Mar.

plus one extra treatment:

EXTRA

NONE No nitrogen fertilizer or inhibitor

NOTE: Dicyandiamide and hydroquinone were applied at 25 and 10 kg respectively in combination with SEEDBD N 0 and at 33 kg and 13 kg with SEEDBED N 50.

Basal applications: Weedkillers: TCA at 11 kg in 250 l. Propyzamide and 3, 6-dichloropicolinic acid (as 'Matrikerb' at 1.6 kg) in 500 l. Fungicide: Prochloraz at 0.50 kg in 250 l. Desiccant: Diquat at 0.6 kg ion with 'Agral', a wetting agent at 0.5 l, in 500 l.

Seed: Jet Neuf, dressed gamma HCH, thiram, fenpropimorph and iprodione, sown at 9.0 kg.

Cultivations, etc.: - Disced: 17 Aug, 1983. TCA applied: 23 Aug. Spring-tine cultivated: 24 Aug. Seed direct drilled: 25 Aug. 'Matrikerb' applied: 25 Oct. Fungicide applied: 18 Apr, 1984. Desiccant applied: 25 July. Combine harvested: 30 July. Previous crops: W. wheat 1982, w. barley 1983.

84/R/RA/2

- NOTES: (1) Dry matter and N contents of plants were measured in February, May and June
 (2) S contents of leaves were measured in May.
 (3) Oil and protein contents of grain were measured.

GRAIN (AT 90% DM) TONNES/HECTARE

***** TABLES OF MEANS *****

SEEDBD N	0	50	MEAN	
N INHIB				
AN O	3.10	3.16	3.13	
PU O	2.87	2.83	2.85	
PU DIC	2.65	2.86	2.76	
PU HYD	3.09	3.04	3.07	
MEAN	2.93	2.97	2.95	
SPRING N	75E+75L	150M	MEAN	
N INHIB				
AN O	3.06	3.21	3.13	
PU O	2.92	2.78	2.85	
PU DIC	2.79	2.72	2.76	
PU HYD	3.06	3.08	3.07	
MEAN	2.96	2.95	2.95	
SPRING N	75E+75L	150M	MEAN	
SEEDBD N				
0	3.01	2.85	2.93	
50	2.90	3.04	2.97	
MEAN	2.96	2.95	2.95	
SEEDBD N	0	50		
SPRING N	75E+75L	150M	75E+75L	150M
N INHIB				
AN O	3.13	3.08	2.98	3.35
PU O	3.00	2.74	2.83	2.82
PU DIC	2.71	2.59	2.87	2.85
PU HYD	3.18	3.01	2.94	3.14
NONE	1.33			
GRAND MEAN	2.86			

84/R/RA/2

GRAIN (AT 90% DM) TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	N INHIB	SEEDBD N	SPRING N	N INHIB SEEDBD N
SED	0.089	0.063	0.063	0.126
TABLE	N INHIB SPRING N	SEEDBD N SPRING N	N INHIB SEEDBD N SPRING N & NONE	
SED	0.126	0.089	0.179	

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	16	0.179	6.3

GRAIN MEAN DM% 87.6

PLOT AREA HARVESTED 0.00465