

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



Potatoes

Rothamsted Research

Rothamsted Research (1985) *Potatoes* ; Yields Of The Field Experiments 1984, pp 326 - 339 - DOI: <https://doi.org/10.23637/ERADOC-1-32>

84/R/P/1

POTATOES

VARIETIES AND STEM CANKER

Object: To study the effects of stem canker (*Rhizoctonia solani*) on plant growth and yield of a range of early and maincrop potato varieties using chitted and unchitted seed - Gt. Knott II.

Sponsors: G.A. Hide, P.J. Read, J.P. Sandison.

Design: Early varieties: 3 randomised blocks of 20 plots.
Maincrop varieties: 3 randomised blocks of 28 plots.

Whole plot dimensions: 3.0 x 10.7.

Treatments:

To EARLY varieties, all combinations of:-

- | | |
|-------------|-------------------------------|
| 1. VARIETY | Varieties: |
| A COMET | Arran Comet |
| ESTIMA | Estima |
| M PEER | Maris Peer |
| U PRINCE | Ulster Prince |
| U SCEPTR | Ulster Sceptre |
| 2. INOCULUM | Inoculum to seed at planting: |
| NONE | None |
| RHIZOCT | R. solani inoculum |
| 3. SD TREAT | Seed treatment: |
| NONE | |
| CHITTED | |

To MAINCROP varieties, all combinations of:-

- | | |
|-------------|-------------------------------|
| 1. VARIETY | Varieties: |
| CARA | Cara |
| DESIREE | Desiree |
| K EDWARD | King Edward |
| M PIPER | Maris Piper |
| P CROWN | Pentland Crown |
| P SQUIRE | Pentland Squire |
| RECORD | Record |
| 2. INOCULUM | Inoculum to seed at planting: |
| NONE | None |
| RHIZOCT | R. solani inoculum |

84/R/P/1

3. SD TREAT Seed treatment:

NONE
CHITTED

NOTE: *Rhizoctonia inoculum* was grown on horticultural vermiculite and sprinkled over seed tubers at planting before covering.

Basal applications: Manures: FYM at 45 t. (10:10:15+4.5 Mg) at 1960 kg.
Weedkillers: Linuron at 1.3 kg with paraquat at 0.50 kg ion in 500 l.
Fungicide: Fentin hydroxide at 0.28 kg in 200 l on seven occasions, applied with the insecticide on the first and third occasion.
Insecticide: Pirimicarb at 0.14 kg on two occasions. Desiccant: Diquat at 0.56 kg ion in 200 l.

Cultivations, etc:- Subsoiled, tines 45 cm deep, 76 cm apart: 6 Oct, 1983.
FYM applied: 4 Nov. Ploughed: 9 Nov. NPK Mg applied, heavy spring-tine cultivated: 23 Mar, 1984. Rotary harrowed: 10 Apr. Early potatoes planted by hand: 11 Apr. Maincrop potatoes planted by hand: 12 Apr. Weedkillers applied: 4 May. Fungicide with insecticide applied: 19 June, 17 July. Fungicide alone applied: 3 July, 30 July, 13 Aug, 28 Aug, 11 Sept. Haulm desiccant applied: 4 Oct. Lifted: 16 Oct. Previous crops: S. barley 1982, w. oats 1983.

NOTE: Plant samples were taken on four occasions to assess stem canker, weight of foliage and weight and numbers of tubers.

84/R/P/1 EARLY POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

INOCULUM VARIETY	NONE	RHIZOCT	MEAN
A COMET	24.3	22.5	23.4
ESTIMA	33.3	30.8	32.1
M PEER	20.3	17.2	18.7
U PRINCE	19.8	16.9	18.3
U SCEPTR	22.1	19.3	20.7

MEAN 24.0 21.3 22.7

SD TREAT VARIETY	NONE	CHITTED	MEAN
A COMET	23.9	23.0	23.4
ESTIMA	32.4	31.7	32.1
M PEER	17.7	19.8	18.7
U PRINCE	18.0	18.6	18.3
U SCEPTR	20.3	21.2	20.7

MEAN 22.5 22.8 22.7

SD TREAT INOCULUM	NONE	CHITTED	MEAN
NONE	23.6	24.3	24.0
RHIZOCT	21.3	21.4	21.3

MEAN 22.5 22.8 22.7

INOCULUM SD TREAT VARIETY	NONE	CHITTED	RHIZOCT NONE	CHITTED
A COMET	24.3	24.4	23.5	21.5
ESTIMA	34.2	32.4	30.7	31.0
M PEER	18.9	21.8	16.5	17.8
U PRINCE	19.3	20.2	16.8	16.9
U SCEPTR	21.5	22.8	19.1	19.5

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	INOCULUM	SD TREAT	VARIETY INOCULUM
SED	1.44	0.91	0.91	2.03

TABLE	VARIETY SD TREAT	INOCULUM SD TREAT	VARIETY INOCULUM SD TREAT
SED	2.03	1.29	2.88

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	38	3.52	15.5

84/R/P/1 EARLY POTATOES

PERCENTAGE WARE 4.44CM (1.75 INCH) RIDDLE

***** TABLES OF MEANS *****

INOCULUM VARIETY	NONE	RHIZOCT	MEAN
A COMET	39.7	65.5	52.6
ESTIMA	77.0	84.0	80.5
M PEER	32.3	50.5	41.4
U PRINCE	65.3	80.3	72.8
U SCEPTR	42.5	67.5	55.0
MEAN	51.4	69.6	60.5

SD TREAT VARIETY	NONE	CHITTED	MEAN
A COMET	50.7	54.5	52.6
ESTIMA	80.9	80.2	80.5
M PEER	41.9	40.9	41.4
U PRINCE	78.1	67.5	72.8
U SCEPTR	53.5	56.5	55.0
MEAN	61.0	59.9	60.5

SD TREAT INOCULUM	NONE	CHITTED	MEAN
NONE	54.6	48.1	51.4
RHIZOCT	67.4	71.7	69.6
MEAN	61.0	59.9	60.5

INOCULUM SD TREAT VARIETY	NONE	CHITTED	RHIZOCT NONE	CHITTED
A COMET	35.1	44.2	66.2	64.7
ESTIMA	79.8	74.3	82.0	86.0
M PEER	39.7	24.9	44.1	56.9
U PRINCE	74.8	55.8	81.4	79.2
U SCEPTR	43.9	41.1	63.1	71.9

PLOT AREA HARVESTED 0.00103

84/R/P/1 MAIN CROP POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

INOCULUM VARIETY	NONE	RHIZOCT	MEAN
CARA	47.7	43.0	45.4
DESIREE	29.3	27.4	28.3
K EDWARD	32.1	25.9	29.0
M PIPER	30.1	28.4	29.2
P CROWN	44.4	39.3	41.9
P SQUIRE	40.4	32.9	36.6
RECORD	25.5	22.2	23.9
MEAN	35.6	31.3	33.5

SD TREAT VARIETY	NONE	CHITTED	MEAN
CARA	43.8	46.9	45.4
DESIREE	29.5	27.2	28.3
K EDWARD	27.9	30.1	29.0
M PIPER	29.3	29.2	29.2
P CROWN	41.6	42.2	41.9
P SQUIRE	37.6	35.7	36.6
RECORD	24.3	23.5	23.9
MEAN	33.4	33.5	33.5

SD TREAT INOCULUM	NONE	CHITTED	MEAN
NONE	35.7	35.6	35.6
RHIZOCT	31.1	31.5	31.3
MEAN	33.4	33.5	33.5

INOCULUM SD TREAT VARIETY	NONE	CHITTED	RHIZOCT NONE	CHITTED
CARA	46.8	48.6	40.8	45.2
DESIREE	30.7	27.9	28.3	26.4
K EDWARD	31.6	32.6	24.2	27.5
M PIPER	29.0	31.2	29.6	27.2
P CROWN	43.9	45.0	39.3	39.4
P SQUIRE	41.9	38.9	33.2	32.5
RECORD	26.3	24.8	22.2	22.2

84/R/P/1 MAIN CROP POTATOES

TOTAL TUBERS TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	INOCULUM	SD TREAT	VARIETY INOCULUM
SED	1.71	0.91	0.91	2.42

TABLE	VARIETY SD TREAT	INOCULUM SD TREAT	VARIETY INOCULUM SD TREAT
SED	2.42	1.29	3.42

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	54	4.19	12.5

84/R/P/1 MAIN CROP POTATOES

PERCENTAGE WARE 4.44CM (1.75 INCH) RIDDLE

***** TABLES OF MEANS *****

INOCULUM VARIETY	NONE	RHIZOCT	MEAN
CARA	90.3	91.4	90.9
DESIREE	77.9	78.5	78.2
K EDWARD	49.9	55.1	52.5
M PIPER	72.3	72.9	72.6
P CROWN	91.2	91.5	91.4
P SQUIRE	90.8	89.1	90.0
RECORD	65.0	68.7	66.8
MEAN	76.8	78.2	77.5

SD TREAT VARIETY	NONE	CHITTED	MEAN
CARA	90.1	91.6	90.9
DESIREE	78.9	77.5	78.2
K EDWARD	51.4	53.5	52.5
M PIPER	70.8	74.4	72.6
P CROWN	91.4	91.3	91.4
P SQUIRE	88.7	91.2	90.0
RECORD	67.8	65.9	66.8
MEAN	77.0	77.9	77.5

SD TREAT INOCULUM	NONE	CHITTED	MEAN
NONE	77.0	76.5	76.8
RHIZOCT	77.0	79.4	78.2
MEAN	77.0	77.9	77.5

INOCULUM SD TREAT VARIETY	NONE	CHITTED	RHIZOCT NONE	CHITTED
CARA	90.0	90.6	90.2	92.6
DESIREE	79.3	76.5	78.5	78.6
K EDWARD	50.0	49.7	52.8	57.3
M PIPER	70.0	74.5	71.5	74.3
P CROWN	91.4	91.0	91.4	91.6
P SQUIRE	89.8	91.9	87.7	90.6
RECORD	68.7	61.3	66.9	70.5

PLOT AREA HARVESTED 0.00103

84/R/P/2

POTATOES

METHODS OF APPLYING FUNGICIDES TO SEED

Object: To compare spraying methods and rates of applying two fungicides to tubers on disease control and yield of potatoes - Gt. Knott II.

Sponsors: G.R. Cayley, G.A. Hide.

Design: 4 randomised blocks of 14 plots.

Whole plot dimensions: 1.5 x 9.52.

Treatments: All combinations of:-

1. FUNGCIDE Fungicides applied to seed tubers:
 IMAZALIL Imazalil
 TOLC MET Tolclofos methyl
2. FUNGRATE Rates of applying fungicides, per tonne of tubers:
 1 5 g imazalil, 12.5 g tolclofos methyl
 2 10 g imazalil, 62.5 g tolclofos methyl
3. FUNGMETH Methods of applying fungicides:
 CNVNTIAL Conventional, hydraulic, sprayer in 2.0 l per tonne of tubers
 SP DS Spinning disc sprayer in 0.8 l per tonne of tubers
 SP DS ES Spinning disc sprayer with electrostatically charged particles in 0.8 l per tonne of tubers

plus one extra treatment:

EXTRA

NONE No fungicides to seed tubers (duplicated)

Basal applications: Manures: FYM at 45 t. (10:10:15+4.5 Mg) at 1960 kg. Weedkiller: Metribuzin at 1.0 kg in 500 l. Fungicide: Fentin hydroxide at 0.28 kg in 200 l applied on seven occasions, the first and third occasion with the insecticide. Insecticide: Pirimicarb at 0.14 kg on two occasions. Desiccant: Diquat at 0.56 kg ion in 200 l.

Seed: King Edward.

Cultivations, etc:- Subsoiled, tines 45 cm deep, 76 cm apart: 6 Oct, 1983. FYM applied: 4 Nov. Ploughed: 9 Nov. NPK Mg applied, heavy spring-tine cultivated: 23 Mar, 1984. Rotary harrowed: 14 Apr. Planted by hand: 16 Apr. Weedkiller applied: 31 May. Fungicide with insecticide applied: 19 June, 17 July. Fungicide alone applied: 3 July, 30 July, 13 Aug, 28 Aug, 11 Sept. Haulm mechanically destroyed, haulm desiccant applied: 4 Oct. Lifted: 16 Oct. Previous crops: S. barley 1982, w. oats 1983.

NOTE: Assessments of stem base infections were made in mid-July.

84/R/P/2

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGRATE	1	2	MEAN		
FUNGCIDE					
IMAZALIL	30.5	32.8	31.7		
TOLC MET	29.9	29.4	29.7		
MEAN	30.2	31.1	30.7		
FUNGMETH	CNVNTIAL	SP DS	SP DS ES	MEAN	
FUNGCIDE					
IMAZALIL	31.5	30.5	33.0	31.7	
TOLC MET	28.9	31.5	28.5	29.7	
MEAN	30.2	31.0	30.8	30.7	
FUNGMETH	CNVNTIAL	SP DS	SP DS ES	MEAN	
FUNGRATE					
1	28.8	31.1	30.9	30.2	
2	31.6	30.9	30.7	31.1	
MEAN	30.2	31.0	30.8	30.7	
FUNGRATE	1			2	
FUNGMETH	CNVNTIAL	SP DS	SP DS ES	CNVNTIAL	SP DS SP DS ES
FUNGCIDE					
IMAZALIL	29.5	29.6	32.6	33.5	31.4 33.4
TOLC MET	28.0	32.6	29.1	29.8	30.4 28.0
NONE	30.7				
GRAND MEAN	30.7				

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	FUNGCIDE	FUNGRATE	FUNGMETH	FUNGCIDE FUNGRATE
SED	0.98	0.98	1.21	1.39
TABLE	FUNGCIDE FUNGMETH	FUNGRATE FUNGMETH	FUNGCIDE FUNGRATE FUNGMETH	
SED	1.70	1.70	2.41	

SED FOR COMPARING NONE WITH ANY ITEM IN FUNGCIDE.FUNGRATE.FUNGMETH TABLE IS 2.09

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	40	3.41	11.1

84/R/P/2

PERCENTAGE WARE 4.44 CM (1.75 INCH) RIDDLE

***** TABLES OF MEANS *****

FUNGRATE	1	2	MEAN			
FUNGCIDE						
IMAZALIL	65.2	65.3	65.3			
TOLC MET	65.9	65.3	65.6			
MEAN	65.6	65.3	65.4			
FUNGMETH	CNVNTIAL	SP DS	SP DS ES	MEAN		
FUNGCIDE						
IMAZALIL	64.0	65.2	66.6		65.3	
TOLC MET	66.6	66.4	63.8		65.6	
MEAN	65.3	65.8	65.2		65.4	
FUNGMETH	CNVNTIAL	SP DS	SP DS ES	MEAN		
FUNGRATE						
1	66.4	65.2	65.1		65.6	
2	64.2	66.4	65.3		65.3	
MEAN	65.3	65.8	65.2		65.4	
FUNGRATE	1				2	
FUNGMETH	CNVNTIAL	SP DS	SP DS ES	CNVNTIAL	SP DS	SP DS ES
FUNGCIDE						
IMAZALIL	64.3	64.7	66.6		63.7	65.7 66.6
TOLC MET	68.5	65.6	63.7		64.7	67.2 64.0
NONE	66.5					
GRAND MEAN	65.6					
PLOT AREA HARVESTED	0.00120					

84/R/P/3

POTATOES

SEED HEALTH

Object: To study the effects of three amounts of pests and disease control on two potato varieties grown for seed - Summerdells I.

Sponsors: R.W. Gibson, R. Harrington, G.A. Hide, G.R. Cayley, D.H. Lapwood.

Design: 2 randomised blocks of 2 plots split into 6.

Whole plot dimensions: 18.0 x 7.62.

Treatments: All combinations of:-

Whole plots

1. VARIETY Varieties:

K EDWARD	King Edward
M PIPER	Maris Piper

Sub plots

2. PATHCONT Pest and pathogen control (in addition to basals):

STANDARD	None
ENHANCED	Seed treatment with tolclofos methyl at 0.24 kg and imazalil at 0.010 kg per tonne of tubers, applied by hydraulic and uncharged electrostatic sprayers respectively. Cypermethrin at 0.04 kg with 7.0 l oil in 500 l applied by hydraulic sprayer on 31 May, 1984. Plants with 'virus' symptoms were removed on 11 June, 20 June and 5 July.

FULL	As for ENHANCED plus:- The imazalil was applied by charged electrostatic sprayer. Plants with 'blackleg' symptoms were removed on 11 June, 20 June and 5 July. Permethrin at 0.10 kg with oil at 7.0 l, in 200 l for the first and last occasions and 500 l for the remainder, was applied on 15 June, 29 June, 16 July, 27 July and (to HAULM D LATER plots only) 13 Aug.
------	---

3. HAULM D Dates of destroying haulm and of lifting:

EARLY	Haulm mechanically destroyed, 3 Aug, 1984. Haulm desiccant applied 6 Aug and potatoes lifted 6 Sept.
LATER	Haulm mechanically destroyed 8 Sept. Haulm desiccant applied 22 Sept and potatoes lifted 11 Oct.

84/R/P/3

Basal applications: Manures: (0:18:36) at 690 kg. (10:10:15+4.5 Mg) at 1960 kg. Weedkillers: Paraquat at 0.80 kg ion in 250 l on two occasions, with the linuron on the second. Linuron at 1.3 kg. Fungicide: Fentin hydroxide at 0.28 kg in 200 l on six occasions (on the fifth and sixth to HAULM D LATER only) applied with the pirimicarb on all but the fifth occasion. Insecticides: Pirimicarb at 0.14 kg on five occasions (on the fifth to HAULM D LATER only). Phorate at 1.7 kg. Haulm desiccant: BOV at 170 l.

Cultivations, etc:- Discd: 2 Sept, 1983. PK applied: 5 Sept. Paraquat applied: 23 Sept. Ploughed: 22 Dec. NPK Mg applied: 3 Apr, 1984. Rotary harrowed, potatoes planted, phorate applied: 10 Apr. Linuron with paraquat applied: 3 May. Fentin hydroxide with pirimicarb applied to all plots: 19 June, 3 July, 17 July, 30 July. Fentin hydroxide applied to HAULM D LATER plots: 13 Aug. Fentin hydroxide with pirimicarb applied to HAULM D LATER plots: 28 Aug. Previous crops: S. barley 1982 and 1983.

NOTE: Aphids were counted throughout the season. Virus and blackleg counts were made in mid-June and mid-July.

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

PATHCONT VARIETY	STANDARD	ENHANCED	FULL	MEAN
K EDWARD	33.8	33.7	25.3	30.9
M PIPER	32.3	31.4	23.8	29.2
MEAN	33.0	32.5	24.6	30.0
HAULM D VARIETY	EARLY	LATER	MEAN	
K EDWARD	24.0	37.8	30.9	
M PIPER	22.2	36.2	29.2	
MEAN	23.1	37.0	30.0	
HAULM D PATHCONT	EARLY	LATER	MEAN	
STANDARD	25.5	40.6	33.0	
ENHANCED	25.1	39.9	32.5	
FULL	18.7	30.4	24.6	
MEAN	23.1	37.0	30.0	

84/R/P/3

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

PATHCONT HAULM D VARIETY	STANDARD EARLY	LATER	ENHANCED EARLY	LATER	FULL EARLY	LATER
K EDWARD	26.3	41.3	26.5	40.8	19.3	31.2
M PIPER	24.7	39.9	23.8	39.1	18.0	29.6

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PATHCONT	HAULM D	VARIETY* PATHCONT
SED	1.51	1.23	2.14

TABLE	VARIETY* HAULM D	PATHCONT HAULM D	VARIETY* PATHCONT HAULM D
SED	1.75	2.14	3.02

* WITHIN THE SAME LEVEL OF VARIETY ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	10	3.02	10.1

84/R/P/3

PERCENTAGE WARE 4.44CM (1.75 INCH) RIDDLE

***** TABLES OF MEANS *****

PATHCONT VARIETY	STANDARD	ENHANCED	FULL	MEAN
K EDWARD	65.8	59.6	57.0	60.8
M PIPER	51.8	48.5	35.9	45.4
MEAN	58.8	54.1	46.4	53.1

HAULM D VARIETY	EARLY	LATER	MEAN
K EDWARD	52.1	69.5	60.8
M PIPER	34.4	56.5	45.4
MEAN	43.2	63.0	53.1

HAULM D PATHCONT	EARLY	LATER	MEAN
STANDARD	49.0	68.6	58.8
ENHANCED	41.3	66.8	54.1
FULL	39.4	53.5	46.4
MEAN	43.2	63.0	53.1

PATHCONT HAULM D VARIETY	STANDARD EARLY	LATER	ENHANCED EARLY	LATER	FULL EARLY	LATER
K EDWARD	57.5	74.1	51.4	67.8	47.4	66.5
M PIPER	40.5	63.1	31.3	65.8	31.4	40.5

SUB PLOT AREA HARVESTED 0.00457