

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1984

[Full Table of Content](#)



84/R/RN/17 Rates of P and K to the Subsoil - S. Beans. W. Wheat, Potatoes, S. Barley

Rothamsted Research

Rothamsted Research (1985) *84/R/RN/17 Rates of P and K to the Subsoil - S. Beans. W. Wheat, Potatoes, S. Barley*; Yields Of The Field Experiments 1984, pp 96 - 102 - DOI:

<https://doi.org/10.23637/ERADOC-1-32>

84/R/RN/17

RATES OF P AND K TO THE SUBSOIL

Object: To study the effects of a range of rates and frequencies of application of P and K to the subsoil, singly and together, on the yields and nutrient uptakes of a rotation of crops - Meadow.

Sponsors: J. McEwen, A.E. Johnston.

The fourth year, potatoes, s. barley, s. beans, w. wheat.

For previous years see 81-83/R/RN/17.

Design: 4 series (for crops) each of 40 plots.

Whole plot dimensions: 3.0 x 14.0.

Treatments to each series:

TREATMNT Extra P and K and primary cultivation tool in autumn 1980 only except on R plots, treatments repeated each autumn:

	P ₂ O ₅ (kg)	K ₂ O(kg)	Tool	
- - -	0	0	Plough	(duplicated)
P6 K6 T	1000	500 to topsoil	"	(")
- - S	0	0 " "	Wye double-digger	(four plots)
- - SR	0	0 " "	" " "	(duplicated)
P2 - SR	63	0 to subsoil	" " "	
P3 - S	125	0 " "	" " "	
P4 - S	250	0 " "	" " "	
P5 - S	500	0 " "	" " "	(duplicated)
P6 - S	1000	0 " "	" " "	
- K2 SR	0	31 " "	" " "	
- K3 S	0	63 " "	" " "	
- K4 S	0	125 " "	" " "	
- K5 S	0	250 " "	" " "	(duplicated)
- K6 S	0	350 " "	" " "	
P1 K1 SR	31	16 " "	" " "	
P1 K3 SR	31	63 " "	" " "	
P2 K2 SR	63	31 " "	" " "	
P3 K1 SR	125	16 " "	" " "	
P3 K3 SR	125	63 " "	" " "	
P3 K4 S	125	125 " "	" " "	
P4 K3 S	250	63 " "	" " "	
P4 K4 S	250	125 " "	" " "	
P4 K5 S	250	250 " "	" " "	(duplicated)
P4 K6 S	250	350 " "	" " "	
P5 K4 S	500	125 " "	" " "	(duplicated)
P5 K5 S	500	250 " "	" " "	
P5 K6 S	500	350 " "	" " "	
P6 K4 S	1000	125 " "	" " "	
P6 K5 S	1000	250 " "	" " "	
P6 K6 S	1000	350 " "	" " "	

84/R/RN/17

- NOTES: (1) Subsoiling was done with the Wye double-digger which turns a furrow with a conventional plough share, to a depth of 23 cm, and at the same time rotary cultivates the bottom of the adjacent furrow to a further depth of 15 cm. When applying P and K this was distributed ahead of the rotary cultivator.
- (2) The topsoil PK dressing was equally divided before and after ploughing.
- (3) All plots other than R were conventionally ploughed in autumn 1981, 1982 and 1983.
- (4) The rate of 350 kg K_2O applied was in error for 500 kg K_2O .

Standard applications:

- Potatoes: Manures: (10:10:15 + 4.5 Mg) at 1960 kg. Weedkillers: Paraquat at 0.50 kg ion with linuron at 1.3 kg in 500 l. Glyphosate at 1.4 kg in 250 l. Fungicides: Fentin hydroxide at 0.28 kg in 200 l on seven occasions, with the insecticide on the first, third, fourth and sixth occasions. Insecticide: Pirimicarb at 0.14 kg on four occasions. Haulm desiccant: Diquat at 0.56 kg ion in 200 l.
- S. barley: Manures: (20:10:10) at 630 kg. Weedkillers: 3, 6-dichloropicolinic acid at 0.07 kg and bromoxynil at 0.34 kg with mecoprop (as 'CMPP' at 4.2 l) applied with the fungicide in 250 l. Fungicide: Tridemorph at 0.52 kg.
- S. beans: Weedkillers: Glyphosate at 1.4 kg in 250 l. Simazine at 1.2 l in 250 l. Insecticide: Phorate at 5.6 kg.
- W. wheat: Manures: (0:18:36) at 350 kg. 'Nitro-Chalk' at 750 kg. Weedkillers: Mecoprop at 2.0 kg, ioxynil at 0.25 kg and bromoxynil at 0.25 kg in 200 l. Fungicide: Propiconazole at 0.25 kg in 500 l. Insecticide: Pirimicarb at 0.14 kg in 250 l.

- Seed: Potatoes: Pentland Crown.
- S. barley: Triumph, dressed with triadimenol and fuberidazole, sown at 160 kg.
- S. beans: Minden, sown at 240 kg.
- W. wheat: Avalon, sown at 190 kg.

Cultivations, etc.:-

- All crops: Treatments applied by double digger: 7-10 Nov, 1983. Ploughed: 11 Nov.
- Potatoes: Glyphosate applied: 6 Oct, 1983. Heavy spring-tine cultivated twice: 16 Jan, 1984 and a third time: 14 Feb. NPK Mg applied: 3 Apr. Spiked rotary cultivated, potatoes planted: 4 Apr. Rotary ridged: 6 Apr. Linuron and paraquat applied: 3 May. Fentin hydroxide with the insecticide applied: 19 June, 17 July, 30 July, 28 Aug. Fentin hydroxide applied: 3 July, 13 Aug, 11 Sept. Haulm mechanically destroyed: 3 Oct. Desiccant applied: 4 Oct. Lifted: 24 Oct.
- S. barley: Spring-tine cultivated: 14 Nov, 1983. Heavy spring-tine cultivated: 14 Feb, 1984. NPK applied: 7 Mar. Spring-tine cultivated, rotary harrowed, seed sown: 8 Mar. Weedkillers and fungicide applied: 23 May. Combine harvested: 17 Aug.
- S. beans: Glyphosate applied: 6 Oct, 1983. Heavy spring-tine cultivated twice: 16 Jan, 1984, and a third time: 14 Feb. Insecticide applied, heavy spring-tine cultivated, rotary harrowed, seed sown: 20 Mar. Simazine applied: 22 Mar. Combine harvested: 31 Aug.
- W. wheat: Glyphosate applied: 6 Oct, 1983. Spring-tine cultivated, PK applied, spring-tine cultivated, rotary harrowed, seed sown: 14 Nov. N applied: 9 Apr, 1984. Weedkillers applied: 19 Apr. Fungicide applied: 14 June. Insecticide applied: 26 June. Combine harvested: 22 Aug.

84/R/RN/17

SERIES II POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	57.7
P6 K6 T	66.1
- - S	61.3
- - SR	56.8
P2 - SR	53.1
P3 - S	52.3
P4 - S	59.0
P5 - S	63.0
P6 - S	57.7
- K2 SR	60.1
- K3 S	62.4
- K4 S	61.3
- K5 S	64.1
- K6 S	58.1
P1 K1 SR	52.4
P1 K3 SR	62.3
P2 K2 SR	55.7
P3 K1 SR	61.5
P3 K3 SR	60.8
P3 K4 S	58.3
P4 K3 S	58.9
P4 K4 S	59.3
P4 K5 S	65.1
P4 K6 S	61.8
P5 K4 S	63.1
P5 K5 S	65.6
P5 K6 S	64.2
P6 K4 S	62.0
P6 K5 S	60.9
P6 K6 S	65.6
MEAN	60.8

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
SED	2.91 MIN REP 2.30 MAX-MIN

* SED APPLIES ONLY TO - - -, P6 K6 T, - - S, - - SR, P5 - S, - K5 S, P4 K5 S AND P5 K4 S

TREATMNT
MAX-MIN - - S V ANY OF REMAINDER
MIN REP ANY OF THE REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	2.06	3.4

84/R/RN/17

SERIES II POTATOES

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

TREATMNT	
- - -	98.0
P6 K6 T	98.4
- - S	98.0
- - SR	98.0
P2 - SR	98.1
P3 - S	98.3
P4 - S	98.8
P5 - S	97.6
P6 - S	98.2
- K2 SR	97.7
- K3 S	98.2
- K4 S	98.6
- K5 S	98.7
- K6 S	97.7
P1 K1 SR	99.2
P1 K3 SR	98.8
P2 K2 SR	98.9
P3 K1 SR	98.3
P3 K3 SR	98.8
P3 K4 S	99.1
P4 K3 S	98.7
P4 K4 S	98.0
P4 K5 S	98.0
P4 K6 S	98.1
P5 K4 S	98.2
P5 K5 S	98.4
P5 K6 S	98.5
P6 K4 S	97.6
P6 K5 S	98.4
P6 K6 S	97.7
MEAN	98.2

PLOT AREA HARVESTED 0.00210

84/R/RN/17

SERIES III SPRING BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	9.22
P6 K6 T	9.37
- - S	9.17
- - SR	9.13
P2 - SR	9.22
P3 - S	9.31
P4 - S	9.47
P5 - S	9.21
P6 - S	9.71
- K2 SR	9.01
- K3 S	9.51
- K4 S	9.01
- K5 S	9.20
- K6 S	8.72
P1 K1 SR	9.06
P1 K3 SR	9.37
P2 K2 SR	9.17
P3 K1 SR	9.33
P3 K3 SR	9.37
P3 K4 S	8.80
P4 K3 S	9.19
P4 K4 S	9.41
P4 K5 S	9.41
P4 K6 S	9.54
P5 K4 S	9.15
P5 K5 S	9.45
P5 K6 S	9.37
P6 K4 S	9.25
P6 K5 S	9.40
P6 K6 S	9.35
MEAN	9.25

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
-----	-----
SED	0.196 MIN REP
	0.155 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.139	1.5

GRAIN MEAN DM% 85.7

PLOT AREA HARVESTED 0.00286

84/R/RN/17

SERIES IV SPRING BEANS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	4.50
P6 K6 T	4.06
- - S	4.14
- - SR	4.42
P2 - SR	4.73
P3 - S	4.89
P4 - S	4.57
P5 - S	4.61
P6 - S	3.23
- K2 SR	6.06
- K3 S	4.15
- K4 S	4.31
- K5 S	4.25
- K6 S	4.86
P1 K1 SR	4.67
P1 K3 SR	4.16
P2 K2 SR	4.80
P3 K1 SR	5.22
P3 K3 SR	4.41
P3 K4 S	5.00
P4 K3 S	3.78
P4 K4 S	3.62
P4 K5 S	4.62
P4 K6 S	4.41
P5 K4 S	3.89
P5 K5 S	3.74
P5 K6 S	4.04
P6 K4 S	4.04
P6 K5 S	3.88
P6 K6 S	3.95
MEAN	4.34

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT*
SED	0.453 MIN REP
	0.358 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.320	7.4
GRAIN MEAN DM%	88.0		
PLOT AREA HARVESTED	0.00386		

84/R/RN/17

SERIES I WINTER WHEAT

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TREATMNT	
- - -	9.84
P6 K6 T	10.34
- - S	9.73
- - SR	10.56
P2 - SR	11.19
P3 - S	11.92
P4 - S	10.66
P5 - S	10.48
P6 - S	10.94
- K2 SR	11.02
- K3 S	10.32
- K4 S	10.13
- K5 S	9.97
- K6 S	10.44
P1 K1 SR	10.06
P1 K3 SR	10.36
P2 K2 SR	10.04
P3 K1 SR	10.37
P3 K3 SR	11.49
P3 K4 S	9.75
P4 K3 S	10.64
P4 K4 S	10.36
P4 K5 S	10.26
P4 K6 S	10.24
P5 K4 S	10.76
P5 K5 S	10.04
P5 K6 S	10.40
P6 K4 S	10.03
P6 K5 S	9.08
P6 K6 S	10.08
MEAN	10.32

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TREATMNT
-----	-----
SED	0.449 MIN REP
	0.355 MAX-MIN

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	10	0.317	3.1

GRAIN MEAN DM% 86.7

PLOT AREA HARVESTED 0.00286