

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1985

[Full Table of Content](#)



## 85/W/CS/99 Effects of Breaks on Take-all - W. Oats

### Rothamsted Research

Rothamsted Research (1986) *85/W/CS/99 Effects of Breaks on Take-all - W. Oats* ; Yields Of The Field Experiments 1985, pp 115 - 115 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/W/CS/99

EFFECTS OF BREAKS ON TAKE-ALL

Object: To study factors affecting the incidence of take-all (*Gaeumannomyces graminis*) and their effects on yields of s. cereals - Woburn, Butt Furlong.

Sponsor: D. Hornby.

The 14th year, w. oats.

For previous years see 72/W/CS/99(t) and 73-84/W/CS/99.

Design: 2 randomised blocks of 9 plots, 6 of which are split into 2.

Whole plot dimensions: 5.34 x 15.2.

Note: Yields were not taken and no new treatments were applied.

Standard applications: Manures: (5:14:30) at 340 kg. N at 120 kg as 'Nitro-Chalk' (27.5% N). Weedkillers: Methabenzthiazuron at 1.6 kg in 250 l. Clopyralid at 0.07 kg with bromoxynil octanoate at 0.34 kg and mecoprop at 2.5 kg in 250 l with the growth regulator. Growth regulator: Chlormequat on two occasions, on the first occasion (as 'Power 3 C' at 2.0 l) with the weedkillers, on the second occasion (as 'Power 3 C' at 4.2 l) with the fungicide. Fungicide: Tridemorph at 0.52 kg in 250 l.

Seed: Panema, sown at 160 kg.

Cultivations, etc.: - Discd: 18 Sept, 1984. Subsoiled, tines 142 cm apart, 56 cm deep, ploughed: 6 Oct. NPK applied: 15 Oct. Rotary harrowed, seed sown: 16 Oct. Methabenzthiazuron applied: 27 Oct. N applied: 16 Apr, 1985. Remaining weedkillers and growth regulator applied: 17 Apr. Fungicide and growth regulator applied: 26 May. Combine harvested: 22 Aug.