

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1985

[Full Table of Content](#)



## 85/R/CS/212 Seasonal Effects of Take-all - W. Wheat, S. Beans

### Rothamsted Research

Rothamsted Research (1986) *85/R/CS/212 Seasonal Effects of Take-all - W. Wheat, S. Beans* ; Yields Of The Field Experiments 1985, pp 123 - 124 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/CS/212

SEASONAL EFFECTS OF TAKE-ALL

Object: To study the incidence of take-all (*Gaeumannomyces graminis*) in continuous w. wheat and in first and second w. wheats after a break - Great Harpenden I.

Sponsor: D. Hornby.

The eighth year, s. beans, w. wheat.

For previous years see 78-84/R/CS/212.

Design: 3 randomised blocks of 4 plots.

Whole plot dimensions: 5.33 x 31.4.

Treatments:

PREVCROP Previous crops before w. wheat 1985:

	1978	1979	1980	1981	1982	1983	1984
CONT W	W	W	W	W	W	W	W
FIRST W	W	W	BE	W	W	BE	W
BEANS	BE	W	W	BE	W	W	BE

BE = s. beans, W = w. wheat

NOTE: An additional crop sequence was in s. beans 1985, yields not taken.

Standard applications:

W. wheat: Manures: (0:20:20) at 330 kg, combine drilled.  
'Nitro-Chalk' (27.5% N) at 330 kg. Weedkillers:  
Isoproturon at 2.0 kg, bromoxynil at 0.34 kg, clopyralid at 0.07 kg and mecoprop (as 'CMPP' at 4.2 l) with the fungicides in 500 l. Fungicides: Prochloraz at 0.4 kg and carbendazim at 0.15 kg.

Seed: W. wheat: Avalon, sown at 190 kg.  
S. beans: Minden, sown at 220 kg.

Cultivations, etc.:-

W. wheat: Ploughed: 8 Oct, 1984. Rotary harrowed, seed sown: 15 Oct. N applied: 12 Apr, 1985. Weedkillers and fungicides applied: 16 Apr. Combine harvested: 29 Aug.  
S. beans: Ploughed: 8 Oct, 1984. Heavy spring-tine cultivated: 19 Mar, 1985. Rotary harrowed, seed sown: 20 Mar. Combine harvested: 27 Sept.

NOTE: Take-all was assessed in soil and in w. wheat roots.

85/R/CS/212

WINTER WHEAT

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PREVCROP	CONT W	FIRST W	BEANS	MEAN
	4.45	4.41	6.08	4.98

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	PREVCROP
-----	-----
SED	0.335

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	4	0.410	8.2

GRAIN MEAN DM% 84.7

PLOT AREA HARVESTED 0.00430