

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1985

[Full Table of Content](#)



## 85/R/CS/246 Effects of Subsoiling and Deep P K - S. Barley

### Rothamsted Research

Rothamsted Research (1986) *85/R/CS/246 Effects of Subsoiling and Deep P K - S. Barley* ; Yields Of The Field Experiments 1985, pp 139 - 140 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/CS/246

EFFECTS OF SUBSOILING AND DEEP PK

Object: To study the effects of thorough subsoil disturbance and the incorporation of P and K into the subsoil on soil and crop parameters and on yield of s. barley - Great Field I.

Sponsors: J. McEwen, A.E. Johnston, D.P. Yeoman.

The sixth year, s. barley.

For previous years see 80-84/R/CS/246.

Whole plot dimensions: 4.27 x 17.7.

Design: 2 replicates of 28 plots, fully randomised.

Treatments: All combinations of:-

1. PK SUB            Extra PK and subsoil treatment (applied autumn/winter 1979/80 only):
  - - -            None, mouldboard ploughed (duplicated)
  - - S            Subsoiled
  - P - S            P to subsoil
  - K S            K to subsoil
  - P K S            PK to subsoil
  - P K T            PK to topsoil, mouldboard ploughed
  
2.        N            Nitrogen fertilizer (kg N) as 'Nitro-Chalk' (27.5% N) (cumulative to previous years):
  - 0
  - 40
  - 80
  - 120

- NOTES: (1) Rates of P and K were 1000 kg P205, as superphosphate, 500 kg K2O, as muriate of potash.
- (2) Subsoiling was done with Wye double-digger which turns a furrow with a conventional plough share, to a depth of 23 cm, and at the same time rotary cultivates the bottom of the adjacent furrow to a further depth of 15 cm. When applying P and K this was distributed ahead of the rotary cultivator.
- (3) The topsoil PK dressing was equally divided before and after ploughing.
- (4) All treatments were mouldboard ploughed for 1981 to 1985.

Basal applications: Manures: (0:24:24) at 260 kg. Weedkillers: Clopyralid at 0.05 kg and bromoxynil octanoate at 0.24 kg with mecoprop (as 'CMPP' at 3.0 l) applied with the fungicide in 200 l. Fungicide: Tridemorph at 0.52 kg.

Seed: Triumph, dressed triadimenol and fuberidazole sown at 160 kg.

Cultivations, etc.:- Ploughed: 13 Nov, 1984. PK applied, spring-tine cultivated, N treatments applied, spring-tine cultivated again, seed sown: 19 Mar, 1985. Weedkillers and fungicide applied: 10 May. Combine harvested: 22 Aug.

85/R/CS/246

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	N	0	40	80	120	MEAN
PK SUB						
- - -		3.72	5.74	7.64	7.38	6.12
- - S		3.41	6.35	7.54	8.26	6.39
P - S		4.16	5.19	7.59	7.95	6.22
- K S		3.49	5.69	7.35	7.97	6.12
P K S		4.39	6.22	7.60	8.13	6.58
P K T		4.56	6.05	7.44	7.96	6.50
MEAN		3.92	5.86	7.54	7.86	6.29

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	PK SUB	N	PK SUB	
			N	
SED	0.283		0.567	MIN REP
	0.245	0.214	0.491	MAX-MIN
			0.401	MAX REP

PKSUB  
 MAX REP - - -  
 MAX-MIN - - - V ANY OF REMAINDER  
 MIN REP ANY OF REMAINDER

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
WP	32	0.567	9.0

GRAIN MEAN DM% 81.3

STRAW TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	N	0	40	80	120	MEAN
PK SUB						
- - -		1.38	2.80	4.07	5.06	3.33
- - S		1.10	2.85	4.01	5.20	3.29
P - S		1.67	2.51	4.44	5.26	3.47
- K S		1.13	2.93	4.20	4.51	3.19
P K S		1.74	2.94	4.57	5.80	3.76
P K T		1.63	3.25	4.28	5.57	3.68
MEAN		1.43	2.87	4.23	5.21	3.44

STRAW MEAN DM% 82.1

PLOT AREA HARVESTED 0.00217