

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/CS/302 Eyespot Resistance to Mbc - W. Wheat

Rothamsted Research

Rothamsted Research (1986) *85/R/CS/302 Eyespot Resistance to Mbc - W. Wheat* ; Yields Of The Field Experiments 1985, pp 161 - 162 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/CS/302

EYESPOT RESISTANCE TO MBC

Object: To study the development of resistance to MBC fungicides in eyespot and the ability of resistant strains to survive, spread and infect - Meadow.

Sponsors: G.L. Bateman, B.D.L. Fitt.

The first year, w. wheat.

Design: 2 randomised blocks of 4 plots split into 6.

Whole plot dimensions: 12.0 x 24.0.

Treatments: All combinations of:-

Whole plots

1. FUNGICIDE	Fungicides:
NONE	None
CARB	Carbendazim at 0.25 kg
PRO	Prochloraz at 0.40 kg
CARB+PRO	Carbendazim at 0.15 kg + prochloraz at 0.40 kg

Sub plots

2. EYE INOC	Eyespot inoculum:
NATURAL	Natural background population (duplicated)
W 19R 1S	Inoculated with wheat strains in proportion 19 resistant to one sensitive
W 1R 19S	As above but one resistant to 19 sensitive
R 19R 1S	Inoculated with rye strains, 19 resistant to one sensitive
R 1R 19S	As above but one resistant to 19 sensitive

- NOTES: (1) Fungicide treatments were applied in 200 l on 26 Nov, 1984 repeated on 10 Apr, 1985.
(2) The eyespot inoculum was colonised on oat seed and this was broadcast on 18 Oct, 1984.

Basal applications: Manures: Chalk at 5.0 t. (5:14:30) at 340 kg. 'Nitro-Chalk' (27.5% N) at 720 kg. Weedkillers: Chlortoluron at 3.5 kg in 250 l. Clopyralid at 0.07 kg and bromoxynil octanoate at 0.34 kg with mecoprop (as 'CMPP' at 4.2 l) in 200 l. Insecticide: Pirimicarb at 0.14 kg in 200 l.

Seed: Avalon, sown at 170 kg.

Cultivations, etc.:- Chalk applied, ploughed: 2 Oct, 1984. NPK applied, spring-tine cultivated, rotary harrowed, seed sown, chlortoluron applied: 4 Oct. Remaining weedkillers applied: 10 Apr, 1985. N applied: 16 Apr. Insecticide applied: 10 July. Combine harvested: 29 Aug. Previous crops: S. barley 1983, w. wheat 1984.

NOTE: Eyespot and sharp eyespot were assessed in April and July. Eyespot was characterized according to type and MBC resistance.

85/R/CS/302

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE	NATURAL	W 19R 1S	W 1R 19S	R 19R 1S	R 1R19S	MEAN
NONE	6.75	6.55	5.68	6.20	6.82	6.46
CARB	6.78	6.41	6.72	5.98	6.88	6.59
PRO	6.30	5.80	6.93	6.78	6.82	6.49
CARB+PRO	7.71	7.55	6.80	7.24	7.32	7.39
MEAN	6.89	6.58	6.53	6.55	6.96	6.73

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EYE INOC	FUNGCIDE* EYE INOC
SED	0.255	0.510
	0.221	0.442

MIN REP EYE INOC
 MAX-MIN NATURAL V ANY OF THE REMAINDER
 MIN REP ANY OF THE REMAINDER

* WITHIN THE SAME LEVEL OF FUNGCIDE ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	24	0.510	7.6
GRAIN MEAN DM%	84.0		
PLOT AREA HARVESTED	0.00142		