

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/W/B/1 Autumn Disease Control - W. Barley

Rothamsted Research

Rothamsted Research (1986) *85/W/B/1 Autumn Disease Control - W. Barley* ; Yields Of The Field Experiments 1985, pp 236 - 241 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/W/B/1

WINTER BARLEY

AUTUMN DISEASE CONTROL

Object: To examine the effects of autumn disease control, and interactions with growth regulator, in winter barley grown on contrasting soil types - Woburn, White Horse (light land), Broad Mead I (heavy land).

Sponsor: J.F. Jenkyn.

Design: On each site: 2 replicates of 20 plots, fully randomised.

Whole plot dimensions: 2.75 x 13.0.

Treatments: Duplicates of all combinations of:-

1. AUT FUNG Autumn fungicide:
 - NONE None (organo-Hg to seed)
 - ETHIRIMO Ethirimol to seed (over organo-Hg S.D.)
 - FF 4050 FF 4050 to seed
 - FENPROPI Fenpropimorph spray (organo-Hg to seed) at 0.75 kg in 220 l on 10 Dec, 1984
 - TRIADIME Triadimenol and fuberidazole to seed
2. GR EARLY Growth regulator early:
 - NONE None
 - ME EARLY Mepiquat chloride + 2-Chloroethylphosphonic acid at G.S. 30 (as 'Terpal' at 2.0 l) in 250 l, with a wetting agent ('Citowett' at 0.01 l), on 9 Apr, 1985
3. GR LATE Growth regulator late:
 - NONE None
 - ME LATE Mepiquat chloride + 2-Chloroethylphosphonic acid at G.S. 32 (as 'Terpal' at 2.0 l) in 250 l, with a wetting agent ('Citowett' at 0.01 l), on 24 Apr.

Basal applications: Manures: (5:14:30) at 300 kg. N at 150 kg (White Horse), at 120 kg (Broad Mead I) as 'Nitro-Chalk' (27.5% N).
Magnesian limestone at 7.5 t (Broad Mead I). Weedkillers: Paraquat at 0.30 kg ion in 250 l. Isoproturon at 2.0 kg in 250 l applied with the insecticide; mecoprop at 2.0 kg with cyanazine at 0.30 kg in 250 l applied with the prochloraz. Insecticide: Permethrin at 0.06 kg.
Fungicides: Prochloraz at 0.40 kg. Carbendazim at 0.25 kg in 250 l. Carbendazim at 0.15 kg with tridemorph at 0.38 kg and maneb at 1.6 kg in 250 l. Triadimefon at 0.12 kg and captafol at 1.3 kg in 250 l.
Desiccant: Diquat at 0.56 kg ion in 250 l (Broad Mead I).

Seed: Panda, sown at 300 seeds per square metre (150 kg).

85/W/B/1

Cultivations, etc.:-

NPK applied, heavy spring-tine cultivated (White Horse): 4 Sept, 1984. Straw burnt, disced (Broad Mead I): 6 Sept. NPK applied (Broad Mead I): 12 Sept. Paraquat applied: 18 Sept. Magnesian limestone applied, heavy spring-tine cultivated, seed sown (Broad Mead I): 19 Sept. Ploughed (White Horse): 19 Sept. Rotary harrowed, seed sown (White Horse): 26 Sept. Weedkiller with insecticide applied: 1 Nov. N applied: 4 Apr, 1985. Weedkillers with fungicide applied: 10 Apr. Carbendazim applied: 16 Apr. Carbendazim with tridemorph and maneb applied: 3 May. Triadimefon with captafol applied: 30 May. Desiccant applied (Broad Mead I): 31 July. Combine harvested (Broad Mead I): 5 Aug, (White Horse): 13 Aug. Previous crops: (White Horse): W. wheat 1983, s. barley 1984. (Broad Mead I): W. wheat 1983 and 1984.

- NOTES: (1) Seed emergence counts were made in October.
(2) Assessments of leaf and root diseases were made during the season.
(3) Crop samples were taken in May for counts of shoots, ear numbers and assessment of dry weights and in July for grain and ear numbers.

85/W/B/1 WHITE HORSE

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

| GR EARLY AUT FUNG | NONE | ME EARLY | MEAN |
|----------------------|------|----------|------|
| NONE | 6.89 | 5.99 | 6.44 |
| ETHIRIMO | 6.98 | 6.65 | 6.82 |
| FF 4050 | 7.63 | 6.32 | 6.98 |
| FENPROPI | 7.09 | 6.43 | 6.76 |
| TRIADIME | 7.38 | 6.58 | 6.98 |

MEAN 7.20 6.39 6.79

| GR LATE AUT FUNG | NONE | ME LATE | MEAN |
|---------------------|------|---------|------|
| NONE | 7.14 | 5.74 | 6.44 |
| ETHIRIMO | 7.16 | 6.47 | 6.82 |
| FF 4050 | 7.14 | 6.82 | 6.98 |
| FENPROPI | 6.87 | 6.65 | 6.76 |
| TRIADIME | 7.22 | 6.74 | 6.98 |

MEAN 7.11 6.48 6.79

| GR LATE GR EARLY | NONE | ME LATE | MEAN |
|---------------------|------|---------|------|
| NONE | 7.78 | 6.62 | 7.20 |
| ME EARLY | 6.44 | 6.35 | 6.39 |

MEAN 7.11 6.48 6.79

| GR EARLY GR LATE AUT FUNG | NONE | ME LATE | ME EARLY NONE | ME LATE |
|---------------------------------|------|---------|------------------|---------|
| NONE | 7.90 | 5.89 | 6.39 | 5.59 |
| ETHIRIMO | 7.77 | 6.19 | 6.55 | 6.75 |
| FF 4050 | 8.27 | 6.99 | 6.00 | 6.65 |
| FENPROPI | 6.98 | 7.19 | 6.75 | 6.12 |
| TRIADIME | 7.95 | 6.82 | 6.50 | 6.65 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | AUT FUNG GR EARLY | GR EARLY | GR LATE | AUT FUNG GR EARLY |
|-------|----------------------|----------|---------|----------------------|
| SED | 0.561 | 0.355 | 0.355 | 0.793 |

| TABLE | AUT FUNG GR LATE | GR EARLY GR LATE | AUT FUNG GR EARLY GR LATE |
|-------|---------------------|---------------------|---------------------------------|
| SED | 0.793 | 0.501 | 1.121 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|---------|----|-------|------|
| WP | 20 | 1.121 | 16.5 |

GRAIN MEAN DM% 81.2

85/W/B/1 WHITE HORSE

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

| GR EARLY AUT FUNG | NONE | ME EARLY | MEAN |
|----------------------|------|----------|------|
| NONE | 3.63 | 3.41 | 3.52 |
| ETHIRIMO | 4.03 | 3.62 | 3.82 |
| FF 4050 | 4.29 | 3.52 | 3.90 |
| FENPROPI | 3.70 | 3.41 | 3.55 |
| TRIADIME | 3.63 | 3.37 | 3.50 |
| MEAN | 3.85 | 3.47 | 3.66 |

| GR LATE AUT FUNG | NONE | ME LATE | MEAN |
|---------------------|------|---------|------|
| NONE | 3.81 | 3.23 | 3.52 |
| ETHIRIMO | 3.51 | 4.14 | 3.82 |
| FF 4050 | 4.06 | 3.75 | 3.90 |
| FENPROPI | 3.69 | 3.42 | 3.55 |
| TRIADIME | 3.75 | 3.24 | 3.50 |
| MEAN | 3.76 | 3.56 | 3.66 |

| GR LATE GR EARLY | NONE | ME LATE | MEAN |
|---------------------|------|---------|------|
| NONE | 3.89 | 3.82 | 3.85 |
| ME EARLY | 3.64 | 3.29 | 3.47 |
| MEAN | 3.76 | 3.56 | 3.66 |

| GR EARLY GR LATE AUT FUNG | NONE | ME EARLY LATE | ME EARLY NONE | ME LATE |
|---------------------------------|------|------------------|------------------|---------|
| NONE | 4.11 | 3.14 | 3.50 | 3.32 |
| ETHIRIMO | 3.44 | 4.62 | 3.58 | 3.66 |
| FF 4050 | 4.65 | 3.92 | 3.46 | 3.59 |
| FENPROPI | 3.89 | 3.50 | 3.49 | 3.34 |
| TRIADIME | 3.34 | 3.92 | 4.17 | 2.56 |

STRAW MEAN DM% 80.2

PLOT AREA HARVESTED 0.00248

85/W/B/1 BROAD MEAD I

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

| GR EARLY AUT FUNG | NONE | ME EARLY | MEAN |
|----------------------|------|----------|------|
| NONE | 7.79 | 8.03 | 7.91 |
| ETHIRIMO | 8.16 | 8.49 | 8.33 |
| FF 4050 | 8.52 | 7.62 | 8.07 |
| FENPROPI | 8.23 | 8.27 | 8.25 |
| TRIADIME | 8.20 | 8.68 | 8.44 |

MEAN 8.18 8.22 8.20

| GR LATE AUT FUNG | NONE | ME LATE | MEAN |
|---------------------|------|---------|------|
| NONE | 7.52 | 8.30 | 7.91 |
| ETHIRIMO | 8.03 | 8.63 | 8.33 |
| FF 4050 | 7.93 | 8.22 | 8.07 |
| FENPROPI | 7.78 | 8.71 | 8.25 |
| TRIADIME | 8.30 | 8.58 | 8.44 |

MEAN 7.91 8.49 8.20

| GR LATE GR EARLY | NONE | ME LATE | MEAN |
|---------------------|------|---------|------|
| NONE | 7.83 | 8.53 | 8.18 |
| ME EARLY | 7.99 | 8.45 | 8.22 |

MEAN 7.91 8.49 8.20

| GR EARLY GR LATE AUT FUNG | NONE | ME LATE | ME EARLY NONE | ME LATE |
|---------------------------------|------|---------|------------------|---------|
| NONE | 7.11 | 8.46 | 7.92 | 8.14 |
| ETHIRIMO | 7.89 | 8.43 | 8.16 | 8.82 |
| FF 4050 | 9.07 | 7.98 | 6.78 | 8.46 |
| FENPROPI | 7.44 | 9.01 | 8.12 | 8.41 |
| TRIADIME | 7.63 | 8.77 | 8.96 | 8.40 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | AUT FUNG GR EARLY | GR EARLY | GR LATE | AUT FUNG GR EARLY |
|-------|----------------------|----------|---------|----------------------|
| SED | 0.262 | 0.166 | 0.166 | 0.371 |

| TABLE | AUT FUNG GR LATE | GR EARLY GR LATE | AUT FUNG GR EARLY GR LATE |
|-------|---------------------|---------------------|---------------------------------|
| SED | 0.371 | 0.234 | 0.524 |

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|---------|----|-------|-----|
| WP | 20 | 0.524 | 6.4 |

GRAIN MEAN DM% 74.3

85/W/B/1 BROAD MEAD I

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

| GR EARLY AUT FUNG | NONE | ME EARLY | MEAN |
|----------------------|------|----------|------|
| NONE | 4.12 | 3.87 | 4.00 |
| ETHIRIMO | 3.71 | 3.93 | 3.82 |
| FF 4050 | 4.47 | 3.95 | 4.21 |
| FENPROPI | 4.00 | 3.94 | 3.97 |
| TRIADIME | 4.13 | 4.04 | 4.08 |
| MEAN | 4.09 | 3.95 | 4.02 |

| GR LATE AUT FUNG | NONE | ME LATE | MEAN |
|---------------------|------|---------|------|
| NONE | 3.71 | 4.28 | 4.00 |
| ETHIRIMO | 3.64 | 4.00 | 3.82 |
| FF 4050 | 4.37 | 4.05 | 4.21 |
| FENPROPI | 3.64 | 4.30 | 3.97 |
| TRIADIME | 4.04 | 4.12 | 4.08 |
| MEAN | 3.88 | 4.15 | 4.02 |

| GR LATE GR EARLY | NONE | ME LATE | MEAN |
|---------------------|------|---------|------|
| NONE | 3.84 | 4.33 | 4.09 |
| ME EARLY | 3.92 | 3.97 | 3.95 |
| MEAN | 3.88 | 4.15 | 4.02 |

| GR EARLY GR LATE AUT FUNG | NONE | ME LATE | ME EARLY NONE | ME LATE |
|---------------------------------|------|---------|------------------|---------|
| NONE | 3.67 | 4.57 | 3.75 | 3.99 |
| ETHIRIMO | 3.43 | 3.99 | 3.86 | 4.01 |
| FF 4050 | 4.98 | 3.96 | 3.76 | 4.13 |
| FENPROPI | 3.42 | 4.58 | 3.86 | 4.02 |
| TRIADIME | 3.70 | 4.56 | 4.39 | 3.68 |

STRAW MEAN DM% 76.7

PLOT AREA HARVESTED 0.00248