

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/AG/6 Agdell - W. Wheat

Rothamsted Research

Rothamsted Research (1986) *85/R/AG/6 Agdell - W. Wheat* ; Yields Of The Field Experiments 1985, pp 26 - 29 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/AG/6

AGDELL

Object: To study, by crop yields and soil analyses, the residual values of phosphate and potash applied in the period 1848-1951 and further dressings since 1964.

The 16th year of revised scheme, w. wheat.

For previous years see 'Details' 1967 and 1973, and 74-84/R/AG/6.

Treatments: All combinations of:-

Whole plots

1. OLDRESD Fertilizers and organic manures applied to roots every fourth year, in the period 1848-1948:

NONE	None
PKNAMG	P K Na Mg
NPKNAMGC	N P K Na Mg C

N: 48 kg N as sulphate of ammonia
P: 41 kg P as superphosphate
K: 224 kg K as sulphate of potash
Na: 16 kg Na as sulphate of soda
Mg: 11 kg Mg as sulphate of magnesia
C: Castor meal at 2240 kg supplying about 112 kg N

2. RN CROP Rotation 1848-1951 and crop in 1984 (all w. wheat 1985):

F/WHEAT	With fallow: Roots (turnips or swedes), s. barley, fallow, w. wheat 1848-1951. Wheat in 1984.
L/FALLOW	With legume: Roots, s. barley, legume (clover or beans), w. wheat 1848-1951. Fallow in 1984.

Half plots

3. 1964RESID Residues of 1964 treatments:

P
K

Quarter plots

4. PREVCROP Previous cropping 1958-69 on P-test half plots, 1958-70 on K-test half plots:

ARABLE	Arable or fallow
GRASS	Grass

85/R/AG/6

Sixteenth plots

5. P205 64	K20 64	Rates of 1964 treatments(kg):	
		P205 to P-test half plots	K20 to K-test half plots
0	0		
500	315		
1000	630		
2000	1260		

Thirty second plots

6. To RN CROP F/WHEAT. Residues of P205 applied 1970-72 (kg) and in 1979, 1981 and 1984 (kg):

P205 724

(0)0 None
(375)450 375 total in 1970-72, 150 in 1980, 1981 and 1984

To RN CROP F/WHEAT. Residues of K20 applied 1973-76 (kg) and in 1979, 1981 and 1984 (kg):

K20 764

(0)0 None
(870)900 870 total in 1973-76, 300 in 1980, 1982 and 1984

To RN CROP L/FALLOW. Residues of P205 applied 1970-72 (kg) and in 1980, 1982 and 1985 (kg):

P205 725

(0)0 None
(375)450 375 total in 1970-72, 150 in 1980, 1982 and 1985

To RN CROP L/FALLOW. Residues of K20 applied 1973-76 (kg) and in 1980, 1982 and 1985 (kg):

K20 765

(0)0 None
(870)900 870 total in 1973-76, 300 in 1980, 1982 and 1985

Standard applications: To P-test half plots: K20 at 300 kg. To K-test half plots: P205 at 150 kg.

Basal applications: Manures: 'Nitro-Chalk' (27.5% N) at 870 kg.
Weedkillers: Paraquat at 0.60 kg ion in 250 l. Cyanazine at 0.24 kg with mecoprop at 1.6 kg in 200 l. Fungicides: Propiconazole at 0.25 kg with carbendazim and maneb (as 'Septal' at 2.5 kg) in 200 l.

Seed: Avalon, dressed fonofos, sown at 200 kg.

85/R/AG/6

Cultivations, etc.: - Test and standard P and K applied: 25 July, 1984.
 Paraquat applied: 11 Oct. Cultivated by rotary digger: 15 Oct.
 Rotary harrowed, seed sown: 6 Feb, 1985. N applied: 15 Apr.
 Cyanazine and mecoprop applied: 16 May. Fungicides applied: 3 July.
 Combine harvested: 7 Sept.

WHEAT AFTER WHEAT P PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	OLDRES	NONE	(375)450	PKNAMG	(375)450	NPKNAMGC	(375)450
	P205 724	(0)0		(0)0		(0)0	
ARABLE	P205 64						
	0	4.31	5.43	6.08	6.32	4.11	4.83
	500	5.23	6.46	6.46	6.87	5.24	6.04
	1000	6.55	7.24	6.35	6.33	3.91	5.58
GRASS	2000	5.11	6.43	6.53	6.39	5.77	5.47
	0	4.25	6.29	2.46	4.78	4.23	4.99
	500	5.66	5.93	4.72	6.23	3.53	5.84
	1000	5.11	6.62	4.47	6.02	5.84	6.30
	2000	6.13	7.14	6.78	7.10	6.70	5.71

GRAIN MEAN DM% 78.4

PLOT AREA HARVESTED (OLDRES NONE) 0.00113
 PLOT AREA HARVESTED (REMAINDER) 0.00127

WHEAT AFTER WHEAT K PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	OLDRES	NONE	(870)900	PKNAMG	(870)900	NPKNAMGC	(870)900
	K20 764	(0)0		(0)0		(0)0	
ARABLE	K20 64						
	0	6.51	6.64	5.83	7.50	6.92	6.27
	315	6.05	7.02	6.91	6.49	5.94	5.87
	630	6.74	6.69	7.05	6.76	6.58	6.36
GRASS	1260	7.02	6.46	7.38	5.87	6.47	6.30
	0	5.58	7.06	6.23	6.58	6.11	6.64
	315	6.88	7.22	7.21	7.16	5.84	7.16
	630	6.82	6.73	5.99	6.60	6.23	6.49
	1260	6.73	7.18	7.72	6.83	6.43	6.41

GRAIN MEAN DM% 78.4

PLOT AREA HARVESTED (OLDRES NONE) 0.00113
 PLOT AREA HARVESTED (REMAINDER) 0.00127

85/R/AG/6

WHEAT AFTER FALLOW P PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	OLDRES D	NONE	PKNAMG		NPKNAMGC		
	P205 725 P205 64	(0)0	(375)450	(0)0	(375)450	(0)0	(375)450
ARABLE	0	6.39	6.89	7.67	7.12	6.80	6.60
	500	6.86	7.18	7.61	7.32	6.06	7.29
	1000	7.48	7.29	7.23	7.45	8.10	7.21
	2000	7.11	7.18	7.76	7.32	6.97	6.88
GRASS	0	7.36	6.75	7.58	7.23	5.20	7.63
	500	7.22	6.40	6.48	7.34	7.03	7.51
	1000	6.68	6.73	6.94	7.62	7.47	7.74
	2000	7.02	6.78	7.01	7.21	7.81	7.39

GRAIN MEAN DM% 78.6

PLOT AREA HARVESTED (OLDRES D NONE) 0.00113
 PLOT AREA HARVESTED (REMAINDER) 0.00127

WHEAT AFTER FALLOW K PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	OLDRES D	NONE	PKNAMG		NPKNAMGC		
	K20 765 K20 64	(0)0	(870)900	(0)0	(870)900	(0)0	(870)900
ARABLE	0	5.09	6.48	8.16	6.23	7.60	7.81
	315	5.54	7.20	7.31	7.15	7.21	7.07
	630	6.84	6.82	7.33	6.50	7.77	7.34
	1260	7.08	7.13	7.05	7.93	7.49	7.12
GRASS	0	7.00	6.43	3.53	6.54	7.21	7.54
	315	6.70	5.11	5.69	6.36	7.31	7.60
	630	6.84	6.87	7.46	5.05	6.97	7.88
	1260	6.88	3.79	7.20	7.27	6.82	7.86

GRAIN MEAN DM% 77.4

PLOT AREA HARVESTED (OLDRES D NONE) 0.00113
 PLOT AREA HARVESTED (REMAINDER) 0.00127