Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1985



Full Table of Content

85/R/BE/4 Varieties - W. Beans

Rothamsted Research

Rothamsted Research (1986) 85/R/BE/4 Varieties - W. Beans; Yields Of The Field Experiments 1985, pp 261 - 262 - DOI: https://doi.org/10.23637/ERADOC-1-19

85/R/BE/4

WINTER BEANS

VARIETIES

Object: To compare agronomic characters and yields of three varieties of w. beans sown at two populations - Long Hoos VI/VII 1.

Sponsors: J. McEwen, D.P. Yeoman.

Design: 3 randomised blocks of 6 plots.

Whole plot dimensions: 2.03 x 2.13.

Treatments: All combinations of:-

1. VARIETY Varieties:

BANNER Banner
BEAGLE Maris Beagle
BOURDON Bourdon

2. POPULATN Plant populations per hectare:

Target population Mean population achieved 130 130,000 120,000 390,000 360,000

NOTE: Seed was sown by hand in rows 51 cm apart.

Basal applications: Manures: Chalk at 2.9 t. Muriate of potash at 520 kg. Weedkillers: Glyphosate at 1.4 kg in 220 l; trietazine at 1.2 kg with simazine at 0.17 kg in 220 l. Fungicides: Benomyl at 0.56 kg plus chlorothalonil at 1.0 kg applied with the second application of deltamethrin; propiconazole at 0.12 kg in 220 l on two occasions. Insecticides: Deltamethrin at 0.0075 kg in 220 l on two occasions; pirimicarb at 0.14 kg in 220 l.

Cultivations, etc.:- Chalk applied: 11 Sept, 1984. K applied: 25 Sept. Glyphosate applied: 26 Sept. Ploughed: 15 Oct. Spring-tine cultivated twice, seed sown, trietazine and simazine applied: 16 Oct. Deltamethrin applied: 2 May, 1985. Deltamethrin with benomyl and chlorothalonil applied: 22 May. Propiconazole applied: 3 June. Pirimicarb applied: 20 June. Propiconazole applied: 9 July. Harvested by hand: 5 Sept. Previous crops: Potatoes 1983, s. barley 1984.

NOTE: Plant counts were made after establishment. Components of yield were measured at maturity. N content of grain was measured. Plant height was measured at the end of May.

85/R/BE/4

GRAIN TONNES/HECTARE

**** TABLES OF MEANS ****

POPULATN VARIETY	130	390	MEAN
BANNER	4.43	5.20	4.81
BEAGLE	5.23	5.03	5.13
BOURDON	5.23	5.48	5.36
MEAN	4.96	5.24	5.10

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	VARIETY	POPULATN	VARIETY POPULATN
SED	0.365	0.298	0.515

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM DF SE CV% BLOCK.WP 10 0.631 12.4

GRAIN MEAN DM% 81.0

PLOT AREA HARVESTED 0.00015