

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/BE/8 Erynia and Aphid Control - S. Beans

Rothamsted Research

Rothamsted Research (1986) *85/R/BE/8 Erynia and Aphid Control - S. Beans ; Yields Of The Field Experiments 1985*, pp 267 - 268 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/BE/8

SPRING BEANS

ERYNIA AND APHID CONTROL

Object: To study the effects of two methods of applying the aphid-pathogenic fungus *Erynia neoaphidis* on the number of black aphids (*Aphis fabae*) and on the yield of s. beans - Little Knott I.

Sponsor: S.K. Mardell.

Design: 5 randomised blocks of 4 plots.

Whole plot dimensions: 2.67 x 2.13.

Treatments:

APH CONT	Chemical and biological aphid control:
NONE	None
EN CULT	<i>E. neoaphidis</i> culture, applied as mycelium
EN APHID	<i>E. neoaphidis</i> applied as mummified aphids
PIRIMICA	Pirimicarb at 0.22 kg in 530 l on 2 July, 1985

NOTES: (1) The culture was applied as 2.23 kg dry mycelium in suspension of water at 970 l on 14 July, 1985. Mummified aphids were applied at 0.5 mg per plant on 1 July, repeated on 8 July.
(2) Basal irrigation was applied as follows (mm water):

4 May	12.5
9 May	12.5
18 May	12.5
3 June	12.5
17 June	12.5
5 July	12.5
14 July	<u>12.5</u>
Total	87.5 mm

Basal applications: Weedkiller: Simazine at 1.2 kg in 200 l.

Seed: Minden, sown at 220 kg.

Cultivations, etc.: - Cultivated by rotary digger: 28 Nov, 1984. Deep spring-tine cultivated, spring-tine cultivated, seed sown: 19 Mar, 1985. Weedkiller applied: 25 Mar. Combine harvested: 27 Sept. Previous crops: S. barley 1983, potatoes 1984.

NOTES: (1) Samples of live aphids were examined for infection with *Erynia* and other fungal pathogens at weekly intervals during June and July.
(2) Aphid numbers were assessed weekly from mid-June to early August.

85/R/BE/8

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

APH CONT	NONE	EN CULT	EN APHID	PIRIMICA	MEAN
	4.52	4.41	4.74	4.87	4.64

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	APH CONT
-----	-----
SED	0.292

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.461	9.9

GRAIN MEAN DM% 89.6

PLOT AREA HARVESTED 0.00024