

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/BE/11 Control of Rust - S. Beans

Rothamsted Research

Rothamsted Research (1986) *85/R/BE/11 Control of Rust - S. Beans ; Yields Of The Field Experiments 1985*, pp 273 - 276 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/BE/11

SPRING BEANS

CONTROL OF RUST

Object: To study the effects of fungicides on the control of rust (*Uromyces viciae-fabae*) and on the yield of unirrigated and irrigated s. beans - Long Hoos VI/VII I.

Sponsors: J. McEwen, D.P. Yeoman.

Design: 2 randomised blocks of 2 plots split into 12.

Whole plot dimensions: 2.03 x 2.13.

Treatments: All combinations of:-

Whole plots

1. IRRIGATN	Irrigation:
0	None
I	Irrigated (33 mm)

Sub plots

2. C S FUNG	Fungicide to control chocolate spot but not rust:
NONE	None
BENOMYL	Benomyl at 0.56 kg in 220 l on 1 July, 1985, 31 July
3. RUSTFUNG	Fungicides to control rust:
MAN+MANC	Maneb at 0.8 kg + mancozeb at 0.8 kg in 340 l
PROPICON	Propiconazole at 0.12 kg in 340 l
4. RFNGTIME	Times of applying fungicides to control rust:
DOUBLE	1 July, 1985, 31 July
QUADRUP	1 July, 15 July, 31 July, 12 Aug

plus two extra sub plot treatments:

EXTRA

NONE	No fungicides (duplicated)
BENOMYL	Benomyl at 0.56 kg in 220 l on 1 July, 31 July (duplicated)

NOTE: Irrigation was applied to lessen a post-flowering soil-moisture deficit from 50 mm to 25 mm. This was necessary on only one occasion, on 18 July, when 33 mm was applied.

Basal applications: Manures: Chalk at 2.9 t. Muriate of potash at 520 kg. Weedkillers: Glyphosate at 1.4 kg in 220 l; trietazine at 1.2 kg with simazine at 0.17 kg and paraquat at 0.40 kg ion in 220 l. Insecticides: Deltamethrin at 0.0075 kg in 220 l on two occasions; pirimicarb at 0.14 kg in 220 l on two occasions.

85/R/BE/11

Seed: Minden, sown at 220 kg.

Cultivations, etc.: - Chalk applied: 11 Sept, 1984. K applied: 25 Sept. Glyphosate applied: 26 Sept. Ploughed: 15 Oct. Spring-tine cultivated, seed sown: 13 Mar, 1985. Trietazine, simazine and paraquat applied: 19 Mar. Deltamethrin applied: 2 May, 23 May. Pirimicarb applied: 20 June, 17 July. Harvested by hand: 18 Sept. Previous crops: Potatoes 1983, s. barley 1984.

NOTE: Plant counts were made after establishment. The incidence of chocolate spot and rust were assessed from early July until maturity. Components of yield were measured at maturity.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

C S FUNG	NONE	BENOMYL	MEAN
IRRIGATN			
0	5.80	5.46	5.63
I	5.95	6.11	6.03
MEAN	5.87	5.78	5.83
RUSTFUNG	MAN+MANC	PROPICON	MEAN
IRRIGATN			
0	6.03	5.24	5.63
I	5.89	6.16	6.03
MEAN	5.96	5.70	5.83
RUSTFUNG	MAN+MANC	PROPICON	MEAN
C S FUNG			
NONE	6.00	5.75	5.87
BENOMYL	5.92	5.65	5.78
MEAN	5.96	5.70	5.83
RFNGTIME	DOUBLE	QUADRU	MEAN
IRRIGATN			
0	5.28	5.99	5.63
I	6.02	6.03	6.03
MEAN	5.65	6.01	5.83
RFNGTIME	DOUBLE	QUADRU	MEAN
C S FUNG			
NONE	5.57	6.18	5.87
BENOMYL	5.73	5.84	5.78
MEAN	5.65	6.01	5.83

85/R/BE/11

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

RFNGTIME	DOUBLE	QUADRUP	MEAN		
RUSTFUNG					
MAN+MANC	5.77	6.15	5.96		
PROPICON	5.53	5.87	5.70		
MEAN	5.65	6.01	5.83		
C S FUNG	NONE		BENOMYL		
RUSTFUNG	MAN+MANC	PROPICON	MAN+MANC	PROPICON	
IRRIGATN					
0	6.11	5.50	5.94	4.98	
I	5.89	6.00	5.89	6.33	
C S FUNG	NONE		BENOMYL		
RFNGTIME	DOUBLE	QUADRUP	DOUBLE	QUADRUP	
IRRIGATN					
0	5.47	6.14	5.08	5.84	
I	5.68	6.22	6.37	5.85	
RUSTFUNG	MAN+MANC		PROPICON		
RFNGTIME	DOUBLE	QUADRUP	DOUBLE	QUADRUP	
IRRIGATN					
0	5.85	6.20	4.70	5.78	
I	5.68	6.10	6.36	5.97	
RUSTFUNG	MAN+MANC		PROPICON		
RFNGTIME	DOUBLE	QUADRUP	DOUBLE	QUADRUP	
C S FUNG					
NONE	5.56	6.44	5.58	5.92	
BENOMYL	5.97	5.86	5.48	5.83	
IRRIGATN	RUSTFUNG	MAN+MANC		PROPICON	
	RFNGTIME	DOUBLE	QUADRUP	DOUBLE	QUADRUP
0	C S FUNG				
	NONE	5.65	6.57	5.28	5.71
	BENOMYL	6.06	5.83	4.11	5.84
I	NONE	5.48	6.31	5.88	6.12
	BENOMYL	5.89	5.89	6.85	5.81
IRRIGATN	0	I	MEAN		
EXTRA					
NONE	4.74	5.10	4.92		
BENOMYL	5.00	5.22	5.11		
MEAN	4.87	5.16	5.02		
GRAND MEAN	5.56				

85/R/BE/11

GRAIN TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	C S FUNG	RUSTFUNG	RFNGTIME
SED	0.265	0.188	0.188	0.188
TABLE	IRRIGATN* C S FUNG	IRRIGATN* RUSTFUNG	C S FUNG RUSTFUNG	IRRIGATN* RFNGTIME
SED	0.265	0.265	0.265	0.265
TABLE	C S FUNG RFNGTIME	RUSTFUNG RFNGTIME	IRRIGATN* EXTRA	IRRIGATN* C S FUNG RUSTFUNG
SED	0.265	0.265	0.375	0.375
TABLE	IRRIGATN* C S FUNG RFNGTIME	IRRIGATN* RUSTFUNG RFNGTIME	C S FUNG RUSTFUNG RFNGTIME	IRRIGATN* C S FUNG RUSTFUNG RFNGTIME
SED	0.377	0.375	0.375	0.531

* WITHIN THE SAME LEVEL OF IRRIGATN ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	26	0.531	9.5

GRAIN MEAN DM% 90.7

PLOT AREA HARVESTED 0.00015