

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/BE/12 Fungicides for Rust Control - S. Beans

Rothamsted Research

Rothamsted Research (1986) *85/R/BE/12 Fungicides for Rust Control - S. Beans* ; Yields Of The Field Experiments 1985, pp 277 - 278 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/BE/12

SPRING BEANS

FUNGICIDES FOR RUST CONTROL

Object: To study the effects of a range of fungicides on the control of rust (*Uromyces viciae-fabae*) and on the yield of s. beans - Long Hoos VI/VII 1.

Sponsors: J. McEwen, D.P. Yeoman.

Design: 2 randomised blocks of 21 plots.

Whole plot dimensions: 2.03 x 2.13.

Treatments: All combinations of:-

1. RUSTFUNG Fungicides to control rust:

BENODANI	Benodanil at 1.1 kg
FENPROP	Fenpropimorph at 0.75 kg
MANEB	Maneb at 0.80 kg
MANCOZEB	Mancozeb at 0.80 kg
MAN+MANC	Maneb at 0.80 kg plus mancozeb at 0.80 kg
PROPICON	Propiconazole at 0.12 kg
THIRAM	Thiram at 2.50 kg
TRIADIME	Triadimefon at 0.50 kg
ZIN+TRID	Zineb polyethylene thiram disulphide at 1.60 kg plus tridemorph at 0.53 kg

 2. RFNGTIME Times of applying fungicides to control rust:

ONCE	Once on 1 July, 1985
TWICE	Twice, on 1 July, 12 Aug
- plus one extra treatment:
- | | |
|-------|---|
| EXTRA | |
| NONE | No fungicides to control rust (triplicated) |

NOTES: (1) All sprays were applied in 220 l.
(2) Irrigation was applied to lessen a post-flowering soil moisture deficit from 50 mm to 25 mm. This was necessary on only one occasion on 18 July, when 33 mm was applied.

Basal applications: Manures: Chalk at 2.9 t. Muriate of potash at 520 kg. Weedkillers: Glyphosate at 1.4 kg in 220 l; trietazine at 1.2 kg with simazine at 0.17 kg and paraquat at 0.40 kg ion in 220 l. Fungicide: Benomyl at 0.56 kg in 220 l. Insecticides: Deltamethrin at 0.0075 kg in 220 l on two occasions; pirimicarb at 0.14 kg in 220 l on two occasions.

Seed: Minden, sown at 220 kg.

85/R/BE/12

Cultivations, etc.:- Chalk applied: 11 Sept, 1984. K applied: 25 Sept. Glyphosate applied: 26 Sept. Ploughed: 15 Oct. Spring-tine cultivated, seed sown: 13 Mar, 1985. Trietazine, simazine and paraquat applied: 19 Mar. Deltamethrin applied: 2 May, 23 May. Pirimicarb applied: 20 June, 17 July. Fungicide applied: 2 July. Harvested by hand: 17 Sept. Previous crops: Potatoes 1983, s. barley 1984.

NOTES: Plant counts were made after establishment. Amounts of chocolate spot and rust were assessed from early July until maturity. Components of yield were measured at maturity.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

RFNGTIME	ONCE	TWICE	MEAN
RUSTFUNG			
BENODANI	6.99	6.99	6.99
FENPROP	6.92	6.80	6.86
MANEB	6.42	5.69	6.06
MANCOZEB	7.30	5.97	6.63
MAN+MANC	7.67	7.17	7.42
PROPICON	5.68	5.65	5.66
THIRAM	6.15	5.94	6.05
TRIADIME	6.51	7.28	6.90
ZIN+TRID	6.91	6.42	6.67
MEAN	6.73	6.43	6.58

EXTRA NONE 5.59

GRAND MEAN 6.44

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	RUSTFUNG	RFNGTIME	RUSTFUNG RFNGTIME
-----	-----	-----	-----
SED	0.524	0.247	0.741

SED FOR COMPARING EXTRA NONE WITH ANY ITEM IN RUSTFUNG.RFNGTIME TABLE IS 0.605

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	22	0.741	11.5

GRAIN MEAN DM% 88.1

PLOT AREA HARVESTED 0.00015