

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1985

[Full Table of Content](#)

Her Majesty's Stationery Office  
LONDON  
1986  
PUBLISHED BY  
HER MAJESTY'S STATIONERY OFFICE  
1986

This report is printed by order of the Secretary of State for the Environment and is available to the public in accordance with the provisions of the Access to Information Act 1982.

Author: Rothamsted  
Publisher: HMSO

## 85/R/LP/2 Desiccants - Lupins

### Rothamsted Research

Rothamsted Research (1986) *85/R/LP/2 Desiccants - Lupins* ; Yields Of The Field Experiments 1985, pp 283 - 284 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/LP/2

LUPINS

DESICCANTS

Object: To study the effects of forms, rates and times of applying desiccants on grain quality and yield of lupins - Long Hoos III 6.

Sponsor: H.L. Jones.

Design: 2 randomised blocks of 14 plots.

Whole plot dimensions: 2.4 x 6.0.

Treatments: All combinations of:-

1. DESICCANT            Forms and rates of desiccants:

DIQUAT 1	Diquat at 0.15 kg ion
DIQUAT 4	Diquat at 0.60 kg ion
NACL 1	Sodium chloride at 6.25 kg
NACL 4	Sodium chloride at 25 kg
GLYPHOS	Glyphosate at 1.1 kg
METAMIT	Metamitron at 2.8 kg

2. DES DATE            Dates of applying desiccants:

4 SEPT	4 Sept, 1985
20 SEPT	20 Sept

plus two extra treatments:

EXTRA

NONE	No desiccant applied
UREA L	Urea at 50 kg on 20 Sept

NOTE: Treatment sprays were applied in 220 l water.

Basal applications: Manures: Chalk at 2.9 t. Muriate of potash at 520 kg. Weedkillers: Glyphosate at 1.4 kg in 220 l; monolinuron at 0.75 kg with paraquat at 0.55 kg ion in 220 l; metamitron at 2.8 kg in 220 l. Fungicide: Benomyl at 0.56 kg in 220 l.

Seed: Vladimir, dressed drazoxolon, sown at 250 kg.

Cultivations, etc.: - Chalk applied: 12 Sept, 1984. Muriate of potash applied: 25 Sept. Glyphosate applied: 26 Sept. Ploughed: 28 Nov. Spring-tine cultivated, seed sown: 2 Apr, 1985. Paraquat and monolinuron applied: 9 Apr. Metamitron applied: 31 May. Fungicide applied: 28 June. Combine harvested: 17 Oct. Previous crops: Sugar beet 1983, s. barley 1984.

NOTE: Observations were made from mid-August on lodging, leaf and pod senescence, stem and pod diseases, plant population, pod numbers per plant, dry matter, shed seed, germinability of seed and seed appearance.

85/R/LP/2

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

DES DATE	4 SEPT	20 SEPT	MEAN
DESICCNT			
DIQUAT 1	3.62	4.28	3.95
DIQUAT 4	3.06	4.01	3.54
NACL 1	3.78	4.14	3.96
NACL 4	3.92	4.04	3.98
GLYPHOS	4.06	3.88	3.97
METAMIT	3.74	4.07	3.90
MEAN	3.70	4.07	3.88

EXTRA	NONE	UREA L	MEAN
	4.08	3.72	3.90

GRAND MEAN 3.89

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	EXTRA	DESICCNT	DES DATE	DESICCNT DES DATE
-----				
SED	0.245	0.173	0.100	0.245

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	13	0.245	6.3

GRAIN MEAN DM% 67.6

PLOT AREA HARVESTED 0.00086