

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/RA/2 Urea and Inhibitors - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1986) *85/R/RA/2 Urea and Inhibitors - W. Oilseed Rape* ; Yields Of The Field Experiments 1985, pp 293 - 295 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/RA/2

WINTER OILSEED RAPE

UREA AND INHIBITORS

Object: To study the effects of adding nitrification inhibitors to prilled urea, applied to the seedbed and in spring on the yield and nitrogen uptake of w. oilseed rape - Black Horse II.

Sponsors: G.A. Rodgers, A. Penny, M.V. Hewitt.

Design: 2 randomised blocks of 18 plots.

Whole plot dimensions: 4.0 x 20.0.

Treatments: All combinations of:-

1. N INHIB Forms of nitrogen and nitrification inhibitor used for seedbed and spring nitrogen applications:

 AN 0 Ammonium nitrate (as 'Nitro-Chalk' (26% N)), no inhibitor
 PU 0 Prilled urea, no inhibitor
 PU DIC Prilled urea and dicyandiamide
 PU HYD Prilled urea and hydroquinone

2. SEEDBD N Nitrogen rates (kg N) to seedbed (on 3 September, 1984):

 0
 50

3. SPRING N Nitrogen rates (kg N) and times in spring:

 75E+75L 75 on 6 Feb, 1985 and 75 on 21 Mar.
 150M 150 on 8 Mar.

plus two extra treatments:

- EXTRA
- SBD ONLY 50 kg N to seedbed only as 'Nitro-Chalk' (26% N), no inhibitor, no N in spring
- NONE No nitrogen fertilizer or inhibitor

NOTE: Dicyandiamide and hydroquinone were applied at 12.5 kg and 10 kg respectively in combination with SEEDBD N 0 and at 18 kg and 13 kg with SEEDBD N 50.

Basal applications: Manures: (0:24:24) at 200 kg. Weedkillers: Propyzamide with clopyralid (as 'Matrikerb' at 1.6 kg) in 500 l; benazolin ethyl ester at 0.30 kg with clopyralid at 0.05 kg in 200 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent ('Agral' at 0.5 l) in 500 l.

Seed: Jet Neuf, seed dressed gamma HCH, thiram and fenpropimorph sown at 8 kg.

85/R/RA/2

Cultivations, etc.:- Disced twice: 31 July, 1984. PK applied: 8 Aug.
 Heavy spring-tine cultivated: 5 Sept. Seed sown: 6 Sept.
 'Matrikerb' applied: 30 Oct. Benazolin ethyl ester with clopyralid
 applied: 6 Mar, 1985. Desiccant applied: 25 July. Combine
 harvested: 12 Aug. Previous crops: W. barley 1983 and 1984.

NOTE: Dry matter and N contents of plants were measured in February, May
 and June. Oil and protein contents of grain were measured.
 Nitrate and ammonium levels in the soil, ammonium losses from main
 dressings and soil pH measurements were taken during the season.
 Disease incidence and severity was assessed once in April.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** TABLES OF MEANS *****

SEEDBD N	0	50	MEAN
N INHIB			
AN O	2.51	2.71	2.61
PU O	2.10	2.19	2.14
PU DIC	2.17	2.34	2.25
PU HYD	2.25	2.52	2.38
MEAN	2.26	2.44	2.35
SPRING N	75E+75L	150M	MEAN
N INHIB			
AN O	2.66	2.56	2.61
PU O	2.36	1.93	2.14
PU DIC	2.33	2.18	2.25
PU HYD	2.58	2.19	2.38
MEAN	2.48	2.21	2.35
SPRING N	75E+75L	150M	MEAN
SEEDBD N			
0	2.44	2.07	2.26
50	2.52	2.36	2.44
MEAN	2.48	2.21	2.35

85/R/RA/2

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** TABLES OF MEANS *****

SEEDBD N	0	150M	50 75E+75L	150M
SPRING N	75E+75L			
N INHIB				
AN 0	2.58	2.43	2.73	2.69
PU 0	2.49	1.70	2.23	2.15
PU DIC	2.25	2.08	2.40	2.28
PU HYD	2.44	2.06	2.72	2.31

EXTRA	SBD ONLY	NONE	MEAN
	1.57	1.27	1.42

GRAND MEAN 2.24

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	N INHIB	SEEDBD N	SPRING N
SED	0.190	0.095	0.067	0.067

TABLE	N INHIB SEEDBD N	N INHIB SPRING N	SEEDBD N SPRING N	N INHIB SEEDBD N SPRING N
SED	0.135	0.135	0.095	0.190

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	17	0.190	8.5

MEAN DM% 82.1

PLOT AREA HARVESTED 0.00472