

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1985

[Full Table of Content](#)



85/R/RA/5 Fungicides and Spray Additive - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1986) *85/R/RA/5 Fungicides and Spray Additive - W. Oilseed Rape* ; Yields Of The Field Experiments 1985, pp 300 - 302 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/RA/5

WINTER OILSEED RAPE

FUNGICIDES AND SPRAY ADDITIVE

Object: To study the effects of times of applying a spray additive and two fungicides, singly and together, on the incidence of diseases and on the yield of w. oilseed rape - Black Horse II.

Sponsor: C.J. Rawlinson.

Design: 2 randomised blocks of 18 plots.

Whole plot dimensions: 3.0 x 12.0.

Treatments: All combinations of:-

1. ADDITIVE Spray additive:
 NONE None
 MIN OIL Mineral oil, 'Actipron' at 5.0 l
2. FUNGICIDE Fungicides, applied at 0.5 kg:
 BENOMYL Benomyl
 PROCHLOR Prochloraz
3. APP TIME Times of applying fungicide:
 AUTUMN Autumn, on 26 Nov, 1984
 SPRING Spring, on 4 Apr, 1985
 AUT+SPNG Autumn and spring as above

plus four extra treatments:

- EXTRA
- | | |
|----------|---|
| NONE | None (triplicated) |
| MINOIL A | Mineral oil as above in autumn |
| MINOIL S | Mineral oil as above in spring |
| MINOILAS | Mineral oil as above in autumn and spring |

NOTE: Fungicide and mineral oil treatments were applied in 200 l of water in autumn and 500 l in spring.

Basal applications: Manures: (0:24:24) at 200 kg. 'Nitro-Chalk' (26% N) at 190 kg followed by 'Nitro-Chalk' (27.5% N) at 900 kg.
Weedkillers: Propyzamide with clopyralid (as 'Matrikerb' at 1.6 kg) in 500 l. Benazolin ethyl ester at 0.30 kg with clopyralid at 0.05 kg in 200 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent ('Agral' at 0.5 l) in 500 l.

Seed: Jet Neuf, seed dressed gamma HCH, thiram and fenpropimorph, sown at 8 kg.

Cultivations, etc.:- Discd twice: 31 July, 1984. PK applied: 8 Aug. First N applied: 10 Aug. Seed sown: 6 Sept. 'Matrikerb' applied: 30 Oct. Second N applied: 27 Feb, 1985. Benazolin ethyl ester with clopyralid applied: 6 Mar. Desiccant applied: 25 July. Combine harvested: 12 Aug. Previous crops: W. barley 1983 and 1984.

85/R/RA/5

NOTE: Establishment counts were made in November. Disease incidence and severity were assessed on five occasions between January and July. Seed shedding on plots was assessed from germinated grain after harvest.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** TABLES OF MEANS *****

FUNGCIDE ADDITIVE	BENOMYL	PROCHLOR	MEAN
NONE	3.50	3.63	3.57
MIN OIL	3.68	3.43	3.56
MEAN	3.59	3.53	3.56

APP TIME ADDITIVE	AUTUMN	SPRING	AUT+SPNG	MEAN
NONE	3.42	3.46	3.82	3.57
MIN OIL	3.47	3.55	3.65	3.56
MEAN	3.45	3.51	3.73	3.56

APP TIME FUNGCIDE	AUTUMN	SPRING	AUT+SPNG	MEAN
BENOMYL	3.56	3.44	3.79	3.59
PROCHLOR	3.33	3.58	3.68	3.53
MEAN	3.45	3.51	3.73	3.56

FUNGCIDE APP TIME ADDITIVE	BENOMYL AUTUMN	PROCHLOR AUTUMN	SPRING	AUT+SPNG	PROCHLOR SPRING	AUT+SPNG
NONE	3.48	3.37	3.29	3.75	3.64	3.89
MIN OIL	3.64	3.30	3.58	3.83	3.52	3.46

EXTRA	NONE	MINOIL A	MINOIL S	MINOILAS	MEAN
	3.62	3.67	3.68	3.24	3.57

GRAND MEAN 3.57

85/R/RA/5

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	EXTRA	ADDITIVE	FUNGCIDE	APP TIME	
SED	0.404 0.297	0.165	0.165	0.202	MIN REP MAX-MIN
TABLE	ADDITIVE FUNGCIDE	ADDITIVE APP TIME	FUNGCIDE APP TIME	ADDITIVE FUNGCIDE APP TIME	
SED	0.233	0.286	0.286	0.404	

EXTRA
MAX-MIN NONE V ANY OF THE REMAINDER
MIN REP ANY OF THE REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	19	0.404	11.3
GRAIN MEAN DM%	86.8		
PLOT AREA HARVESTED	0.00271		