

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1985

[Full Table of Content](#)



## 85/R/RA/6 Growth Regulators - W. Oilseed Rape

### Rothamsted Research

Rothamsted Research (1986) *85/R/RA/6 Growth Regulators - W. Oilseed Rape* ; Yields Of The Field Experiments 1985, pp 303 - 304 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/R/RA/6

WINTER OILSEED RAPE

GROWTH REGULATORS

Object: To study the effects of a range of materials on the control of fungi and on the growth and the yield of w. oilseed rape - Black Horse II.

Sponsors: C.J. Rawlinson, D.P. Yeoman.

Design: 3 randomised blocks of 7 plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments:

CHEMICAL	Chemicals:
NONE	None
BAS11100	'BAS 11100W' at 6.7 l
MEPIQUAT	Mepiquat chloride + ethephon (as 'Terpal' at 3.0 l)
MEP+PROP	Mepiquat chloride + ethephon + propiconazole
PROPICON	Propiconazole at 0.12 kg
PROCHLOR	Prochloraz at 0.40 kg
TRIAPENT	Triapenthenol (as 'UK 140' at 0.7 kg)

NOTES: (1) Treatments were applied in 220 l on 17 Apr, 1985.

(2) Mepiquat chloride + ethephon were applied with a wetting agent ('Citowett' at 0.1 l).

Basal applications: Manures: (0:24:24) at 200 kg. 'Nitro-Chalk' (26% N) at 190 kg followed by 'Nitro-Chalk' (27.5% N) at 900 kg. Weedkillers: Propyzamide with clopyralid (as 'Matrikerb' at 1.6 kg) in 500 l. Benazolin ethyl ester at 0.30 kg with clopyralid at 0.05 kg in 200 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent ('Agral' at 0.5 l) in 500 l.

Seed: Jet Neuf, seed dressed gamma HCH, thiram and fenpropimorph, sown at 8 kg.

Cultivations, etc.: - Discd twice: 31 July, 1984. PK applied: 8 Aug. First N applied: 10 Aug. Seed sown: 6 Sept. 'Matrikerb' applied: 30 Oct. Second N applied: 27 Feb, 1985. Benazolin ethyl ester with clopyralid applied: 6 Mar. Desiccant with wetter applied: 25 July. Combine harvested: 12 Aug. Previous crops: W. barley 1983 and 1984.

NOTE: Disease incidence and severity was assessed on four occasions between April and July. Plant height, internode length, branch number and length, growth stage, flowering and petal size measurements were made in May and July. Lodging was assessed in July. Seed shedding on plots was assessed from germinated grain after harvest.

85/R/RA/6

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CHEMICAL	NONE	BAS11100	MEPIQUAT	MEP+PROP	PROPICON	PROCHLOR	TRIAPENT	ME
	3.00	2.89	2.99	3.49	3.32	3.21	3.28	3.

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	CHEMICAL
-----	-----
SED	0.180

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	12	0.220	6.9

GRAIN MEAN DM% 87.7

PLOT AREA HARVESTED 0.00231