

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1985

[Full Table of Content](#)



## 85/S/RN/2 Rotation II - W. Wheat, W. Beans

### Rothamsted Research

Rothamsted Research (1986) *85/S/RN/2 Rotation II - W. Wheat, W. Beans* ; Yields Of The Field Experiments 1985, pp 41 - 43 - DOI: <https://doi.org/10.23637/ERADOC-1-19>

85/S/RN/2

ROTATION II

Object: To measure, by crop yields and soil analysis, the residual value of P applied as FYM or superphosphate in the periods 1899-1964 and 1965-1967 and of fresh dressings since - Saxmundham.

Sponsor: A.E. Johnston.

The 16th year of revised scheme, w. wheat, w. beans.

For previous years see 'Details' 1967 and 1973, and 74-84/S/RN/2.

Whole plot dimensions: 5.49 x 39.8.

Treatments: From 1899-1964 the experiment tested farmyard manure and nitrogen and phosphate fertilizers applied to a rotation of crops. Since 1965 the treatments have been changed to evaluate old residues of P (from FYM and superphosphate) and new residues from treatments applied 1965-1967. All crops of the rotation - potatoes, s. barley, sugar beet, s. barley - were grown until 1974. The whole experiment was sown to s. barley in 1975 and 1976, alternating w. wheat and s. barley from 1977 to 1979, alternating w. beans and w. wheat in 1980 and 1981, w. wheat alone in 1982 and 1983, w. wheat and w. beans in 1984. Combinations of the following treatments were tested on w. beans and on a first wheat after beans in 1984:

Whole plots

1. RESIDUE

Residues of previous treatments:-

		Approximate total dressing 1899-1964	Total dressing 1965-1967
(0)0	Plot 1	None	None
(D)0	Plot 2	400 tonnes FYM	None
(DP)0	Plot 3	400 tonnes FYM, 2.7 tonnes P205	None
(DP)D2	Plot 4	400 tonnes FYM, 2.7 tonnes P205	100 tonnes FYM
(DP)D2P1	Plot 5	400 tonnes FYM, 2.7 tonnes P205	100 tonnes FYM, 0.56 tonnes P205
(DP)P1	Plot 6	400 tonnes FYM, 2.7 tonnes P205	0.56 tonnes P205
(DP)P2	Plot 7	400 tonnes FYM, 2.7 tonnes P205	1.13 tonnes P205
(DP52)0	Plot 8	326 tonnes FYM, 4.3 tonnes P205 (until 1952 only)	None

85/S/RN/2

Sub plots

2. P Phosphate (total P2O5 applied in each period (kg)):

	1969-71	1973-75	1978*	1980*	1982*	1984*
(0)(0)0	0	0	0	0	0	0
(0)(3)0	0	378	0	0	0	0
(1)(3)1	126	378	120	120	120	120
(2)(3)1	252	378	120	120	120	120
(3)(3)0	378	378	0	0	0	0

\* 1978, 1980, 1982 and 1984 are the years of application for wheat in 1985. Years of application for beans in 1985 were 1979, 1981, 1983 and 1985.

and, for wheat only, some of the combinations of 2 with:-

3. N Nitrogen fertilizer in spring (kg N) as 'Nitro-Chalk' (26% N) in addition to 45 kg N at sowing:

120  
160  
200  
240

NOTE: Plots with the combinations of RESIDUE (DP)D2, (DP)D2P1, (DP)P1, (DP)P2 with P(3)(3)(0) were used for N15 studies, yields not taken.

Standard applications:

W. wheat: Manures: N at 45 kg as prilled urea. Weedkillers: Isoproturon at 2.5 kg in 220 l. Mecoprop at 1.2 kg, bromoxynil at 0.26 kg and ioxynil at 0.26 kg applied with the prochloraz in 220 l. Fungicides: Prochloraz at 0.42 kg. Maneb at 1.6 kg, carbendazim at 0.15 kg and tridemorph at 0.37 kg with captafol at 1.0 kg applied with the insecticide in 220 l. Insecticide: Pirimicarb at 0.14 kg.  
W. beans: Weedkiller: Simazine at 1.1 kg in 220 l. Fungicide: Benomyl at 0.56 kg in 220 l.

Seed: W. wheat: Galahad, sown at 400 seeds per square metre.  
W. beans: Banner, sown at 250 kg.

Cultivations, etc.:-

W. wheat: Ploughed: 12 Oct, 1984. Power harrowed, seed sown, N applied: 30 Oct. Isoproturon applied: 31 Oct. N treatments applied: 17 Apr, 1985. Mecoprop, bromoxynil, ioxynil and prochloraz applied: 23 Apr. Remaining fungicides and insecticide applied: 26 June. Combine harvested: 29 Aug.  
W. beans: P and chalk treatments applied: 6 Sept, 1984. Ploughed: 12 Oct. Power harrowed, seed sown, weedkiller applied: 31 Oct. Fungicide applied: 22 Apr, 1985. Combine harvested (yields not recorded): 19 Sept.

85/S/RN/2

W.WHEAT AFTER BEANS

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

RESIDUE	N P	120	160	200	240
(0)0	(0)(0)0			4.97	6.95
(0)0	(0)(3)0	7.39	5.33		
(0)0	(1)(3)1	8.46		8.61	
(0)0	(2)(3)1		8.51		8.95
(0)0	(3)(3)0		8.89		8.19
(D)0	(0)(0)0	6.19	7.75		
(D)0	(0)(3)0			7.89	7.56
(D)0	(1)(3)1		8.34		8.30
(D)0	(2)(3)1	8.72		7.97	
(D)0	(3)(3)0	7.44		7.52	
(DP)0	(0)(0)0			8.88	8.94
(DP)0	(0)(3)0	7.65	8.54		
(DP)0	(1)(3)1	8.80		8.26	
(DP)0	(2)(3)1		8.88		8.45
(DP)0	(3)(3)0		8.45		8.96
(DP)D2	(0)(0)0	8.34	8.51		
(DP)D2	(0)(3)0			8.36	8.40
(DP)D2	(1)(3)1	9.45		9.05	
(DP)D2	(2)(3)1		9.13		9.49
(DP)D2	(3)(3)0		9.47		9.28
(DP)D2P1	(0)(0)0			7.98	8.79
(DP)D2P1	(0)(3)0	8.49	8.76		
(DP)D2P1	(1)(3)1	9.24		8.07	
(DP)D2P1	(2)(3)1		9.67		9.12
(DP)D2P1	(3)(3)0		10.15		7.97
(DP)P1	(0)(0)0			8.45	8.36
(DP)P1	(0)(3)0	8.22	9.48		
(DP)P1	(1)(3)1		9.41		8.03
(DP)P1	(2)(3)1	8.46		8.16	
(DP)P1	(3)(3)0	8.87		9.56	
(DP)P2	(0)(0)0	9.11	8.95		
(DP)P2	(0)(3)0			9.19	8.15
(DP)P2	(1)(3)1		9.92		8.47
(DP)P2	(2)(3)1	8.59		9.68	
(DP)P2	(3)(3)0	8.85		9.07	
(DP52)0	(0)(0)0	7.68	7.74		
(DP52)0	(0)(3)0			8.61	8.17
(DP52)0	(1)(3)1		8.92		8.84
(DP52)0	(2)(3)1	8.30		8.09	
(DP52)0	(3)(3)0	9.50		8.66	

GRAIN MEAN DM% 83.5

PLOT AREA HARVESTED 0.00073