

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1986

[Full Table of Content](#)



### 86/R/CS/247 Organic Matter and Earthworm Inoculation - S. Barley

#### Rothamsted Research

Rothamsted Research (1987) *86/R/CS/247 Organic Matter and Earthworm Inoculation - S. Barley* ;  
Yields Of The Field Experiments 1986, pp 123 - 123 - DOI:

<https://doi.org/10.23637/ERADOC-1-36>

86/R/CS/247

ORGANIC MATTER AND EARTHWORM INOCULATION

Object: To study the residual effects of inoculating earthworms into arable soil and adding organic materials on the breakdown of straw in the soil by soil animals - Hoosfield.

Sponsor: W. Powell.

The seventh year, s. barley.

For previous years see 80-85/R/CS/247.

Design: 3 randomised blocks of 9 plots.

Whole plot dimensions: 7.85 x 7.62.

Treatments: All combinations of:-

1. WORMINOC(80) Earthworms and inoculation method for 1980 crop only:

NONE	None
EVEN	Earthworms ( <i>Lumbricus terrestris</i> ) applied at 16,700 per hectare in November 1979:
CONC	Evenly spaced throughout
	Concentrated in metre squares, 100 earthworms per square metre
  
2. ORG MATT(85) Forms of organic matter applied until 1985:

NONE	None
STR	Straw at 6.50 t for 1980, 3.25 t for 1981 and 1982
STR+FYM	Straw at 6.50 t for 1980, 3.25 for 1981 and 1982 plus farmyard manure at 40 t for each year including 1985

NOTE: Yields were not taken in 1986.

Basal applications: Manures: 'Nitram' at 380 kg and later at 130 kg.  
Weedkillers: Clopyralid at 0.05 kg and bromoxynil octanoate at 0.24 kg with mecoprop at 2.1 kg applied with the fungicide in 200 l.  
Fungicide: Tridemorph at 0.52 kg.

Seed: Klaxon, sown at 160 kg.

Cultivations, etc.: - Ploughed: 22 Nov, 1985. First N applied: 26 Mar, 1986. Spring-tine cultivated, rotary harrowed, seed sown: 29 Apr. Second N applied: 7 May. Weedkillers and fungicide applied: 2 June. Combine harvested: 6 Sept.