

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1986

[Full Table of Content](#)



### 86/R/CS/302 Eyespot Resistance to Mbc - W. Wheat

#### Rothamsted Research

Rothamsted Research (1987) *86/R/CS/302 Eyespot Resistance to Mbc - W. Wheat* ; Yields Of The Field Experiments 1986, pp 142 - 143 - DOI: <https://doi.org/10.23637/ERADOC-1-36>

86/R/CS/302

EYESPOT RESISTANCE TO MBC

Object: To study the development of resistance to MBC fungicides in eyespot and the ability of resistant strains to survive, spread and infect - Meadow.

Sponsors: G.L. Bateman, B.D.L. Fitt.

The second year, w. wheat.

For previous year see 85/R/CS/302.

Design: 2 randomised blocks of 4 plots split into 6.

Whole plot dimensions: 12.0 x 24.0.

Treatments: All combinations of:-

Whole plots

1. FUNGCIDE Fungicides applied cumulatively to 1985 treatments:

NONE	None
CARB	Carbendazim at 0.25 kg
PRO	Prochloraz at 0.40 kg
CARB+PRO	Carbendazim at 0.15 kg + prochloraz at 0.40 kg

Sub plots

2. EYE INOC Eyespot inoculum, applied in first year only:

NATURAL	Natural background population (duplicated)
W 19R 1S	Inoculated with wheat strains in proportion 19 resistant to one sensitive
W 1R 19S	As above but one resistant to 19 sensitive
R 19R 1S	Inoculated with rye strains, 19 resistant to one sensitive
R 1R 19S	As above but one resistant to 19 sensitive

NOTES: (1) Fungicide treatments were applied in 200 l on 31 Oct, 1985 repeated on 1 May, 1986.

(2) The eyespot inoculum was colonised on oat seed and this was broadcast in October, 1984.

Basal applications: Manures: (5:14:30) at 340 kg. 'Nitram' at 600 kg. Weedkillers: Paraquat at 0.60 kg ion in 200 l. Clopyralid at 0.07 kg, bromoxynil octanoate at 0.34 kg, mecoprop at 2.5 kg and isoproturon at 2.0 kg in 200 l.

Seed: Avalon, sown at 190 kg.

Cultivations, etc.:- Paraquat applied: 23 Sept, 1985. Heavy spring-tine cultivated twice: 26 Sept. NPK applied, disced, rotary harrowed, seed sown: 4 Oct. N applied: 24 Apr, 1986. Weedkillers applied: 25 Apr. Combine harvested: 14 Aug.

NOTE: Eyespot and sharp eyespot were assessed in April and July. Eyespot was characterized according to type and MBC resistance.

86/R/CS/302

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

EYE INOC FUNGICIDE	NATURAL	W 19R 1S	W 1R 19S	R 19R 1S	R 1R 19S	MEAN
NONE	8.79	8.99	8.05	8.67	9.40	8.78
CARB	8.60	8.57	8.35	8.38	8.75	8.54
PRO	8.23	8.26	8.71	8.51	8.44	8.40
CARB+PRO	9.11	8.23	9.13	8.46	9.27	8.89
MEAN	8.68	8.51	8.56	8.50	8.96	8.65

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	EYE INOC	FUNGICIDE* EYE INOC	
SED	0.185	0.370	MIN REP
	0.160	0.321	MAX-MIN

EYE INOC  
MAX-MIN NATURAL V ANY OF THE REMAINDER  
MIN REP ANY OF THE REMAINDER

\* WITHIN THE SAME LEVEL OF FUNGICIDE ONLY

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP.SP	24	0.370	4.3
GRAIN MEAN DM%	84.8		
SUB PLOT AREA HARVESTED	0.00139		