Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1986



Full Table of Content

86/R/CS/318 Nitrophosphates - Potatoes

Rothamsted Research

Rothamsted Research (1987) 86/R/CS/318 Nitrophosphates - Potatoes; Yields Of The Field Experiments 1986, pp 161 - 163 - DOI: https://doi.org/10.23637/ERADOC-1-36

86/R/CS/318

NITROPHOSPHATES

Object: To study the effect of different amounts of water-soluble phosphate in nitrophosphate fertilizers on the growth and P uptake of potatoes following spring barley - Highfield V.

Sponsor: K.H.G. Copestake.

The second year, potatoes.

For previous year see 85/R/B/8.

Design: 3 randomised blocks of 13 plots.

Whole plot dimensions: 3.0 x 21.0.

Treatments: All combinations of:-

1.	P SOL Phosphate water solubility (%):				
	59	Compound fertilizer (16.4 : 14.2 : 17.5) with 59% of the P205 water soluble			
	73	Compound fertilizer (15.9 : 16.2 : 15.3) with 73% of the P205 water soluble			
	95	Compound fertilizer (15.0 : 15.0 : 15.0) with 95% of the P205 water soluble			
2.	P RATE	Rate of phosphate (kg P205) in 1985 and 1986:			
	(20)100 (40)150 (60)200	20 to s. barley in 1985, 100 to potatoes in 1986 40 to s. barley in 1985, 150 to potatoes in 1986 60 to s. barley in 1985, 200 to potatoes in 1986			

plus four extra treatments given rates of compound fertilizer, (3.7: 37.6: 0) with 94% of the P205 water soluble, in 1986 only:

EXTRA(94)

(0)0	None			
(0)100	100	kg	P205	
(0)150	150	kg	P205	
(0)200	200	kg	P205	

NOTE: The compound fertilizers used to apply the phosphate treatments supplied differing amounts of the total 231 kg N and 246.5 kg K20 required on all plots. Additional amounts of N (as 'Nitrotop' 33.5% N) and K20 (as muriate of potash, 60% K20) were applied as needed to achieve this total. Similar balancing was done in the previous year, see 85/R/B/8.

Basal applications: Manures: Kieserite at 500 kg. Weedkiller: Linuron at 1.3 kg in 500 l. Fungicides: Mancozeb at 1.4 kg in 200 l on four occasions, applied with the insecticide on the third and fourth. Fentin hydroxide at 0.28 kg in 200 l on two occasions, applied with the insecticide on the first. Insecticide: Pirimicarb at 0.14 kg on three occasions. Desiccant: Diquat at 0.60 kg ion in 200 l.

Seed: Pentland Crown.

86/R/CS/318

Cultivations, etc.:- Ploughed: 18 Nov, 1985. Spring-tine cultivated: 30 Apr, 1986. Fertilizer treatments applied: 13 May. Mg applied, rotary harrowed, potatoes planted: 14 May. Rotary ridged: 20 May. Weedkiller applied: 30 May. Mancozeb applied: 30 June, 14 July. Mancozeb with pirimicarb applied: 28 July, 12 Aug. Fentin hydroxide with pirimicarb applied: 29 Aug. Fentin hydroxide applied: 11 Sept. Haulm mechanically destroyed: 19 Sept. Desiccant applied: 27 Sept. Lifted: 14 Oct.

NOTE: Emergence and stem counts were made and a mid-season sample was taken for fresh and dry weight. Components of yield were measured. P contents of crop and soil were determined after harvest.

86/R/CS/318

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

P RATE P SOL	(20)100	(40)150	(60)200	MEAN	
59	42.2	43.5	43.7	43.1	
73	40.6	44.3	42.8	42.6	
95	43.6	43.0	46.8	44.5	
MEAN	42.1	43.6	44.4	43.4	
EXTRA(94)	(0)0 38.3	(0)100 40.9	(0)150 42.3	(0)200 45.1	MEAN 41.7

GRAND MEAN 42.9

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	P SOL	P RATE	P SOL P RATE	EXTRA(94)
SED	0.98	0.98	1.97	1.97

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV%

BLOCK.WP 24 2.41 5.6

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

P RATE P SOL	(20)100	(40)150	(60)200	MEAN	
59	98.6	98.6	98.4	98.5	
73	98.9	97.9	98.6	98.4	
95	98.1	98.4	98.4	98.3	
MEAN	98.5	98.3	98.5	98.4	
EXTRA(94)	(0)0 98.6	(0)100 98.2	(0)150 98.2	(0)200 98.0	

GRAND MEAN 98.4

PLOT AREA HARVESTED 0.00165

MEAN 98.3