

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1986

[Full Table of Content](#)



86/R/CS/320 Comparison of Combinable Crops - W. Beans, W. Oats, W. Oilseed Rape, W. Peas, W. Wheat, S. Beans, S. Lupins, S. Peas, Sunflowers and Fallow

Rothamsted Research

Rothamsted Research (1987) *86/R/CS/320 Comparison of Combinable Crops - W. Beans, W. Oats, W. Oilseed Rape, W. Peas, W. Wheat, S. Beans, S. Lupins, S. Peas, Sunflowers and Fallow* ; Yields Of The Field Experiments 1986, pp 166 - 168 - DOI: <https://doi.org/10.23637/ERADOC-1-36>

86/R/CS/320

COMPARISON OF COMBINABLE CROPS

Object: To compare yields and other attributes of a range of combinable crops and to study their effects on a following crop of w. wheat - Long Hoos VI/VII 5.

Sponsors: J. McEwen, D.P. Yeoman, A.E. Johnston.

The first year, w. beans, w. oats, w. oilseed rape, w. peas, w. wheat, s. beans, s. lupins, s. peas, sunflowers and fallow.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 2.5 x 8.0.

Treatments:

CROP	Crops:
W BEANS	W. field beans, <i>Vicia faba</i>
W OATS	W. oats
W RAPE	W. oilseed rape
W PEAS	W. peas, <i>Pisum sativum</i>
W WHEAT	W. wheat
S BEANS	S. field beans, <i>Vicia faba</i>
S LUPINS	S. lupins, <i>Lupinus albus</i>
S PEAS	S. peas, <i>Pisum sativum</i>
SNFLOWER	Sunflower

NOTE: One plot in each block was fallowed.

Standard applications:

All crops and fallow: Manures: Muriate of potash at 520 kg.

Weedkiller: Glyphosate at 0.72 kg in 220 l.

All winter crops only: Insecticide: Deltamethrin at 0.0075 kg in 220 l.

W. beans only: Weedkillers: Simazine at 0.17 kg and trietazine at 1.2 kg with paraquat at 0.28 kg ion in 220 l. Insecticide: Pirimicarb at 0.14 kg in 220 l.

W. oats only: Manures: 'Nitro-Chalk' at 440 kg. Weedkillers: Methabenzthiazuron at 3.1 kg with paraquat at 0.28 kg ion in 220 l.

W. rape only: Manures: 'Nitro-Chalk' at 170 kg and later at 730 kg. Weedkillers: Propyzamide and clopyralid (as 'Matrikerb' at 1.2 kg) in 220 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent ('Cittowet' at 0.10 l) in 220 l.

W. and s. peas only: Insecticide: Pirimicarb at 0.14 kg in 220 l.

W. wheat only: Manures: 'Nitro-Chalk' at 730 kg. Weedkillers: Isoproturon at 2.1 kg with paraquat at 0.28 kg ion in 220 l.

S. lupins only: Fungicide: Benomyl at 0.56 kg in 220 l.

S. beans only: Insecticide: Pirimicarb at 0.14 kg in 220 l.

Sunflowers only: Manures: 'Nitro-Chalk' at 290 kg. Bird repellent: 'Hoppit' at 3.0 l in 220 l.

86/R/CS/320

Seed: W. beans: Banner, sown at 120 kg.
W. oats: Bulwark, sown at 180 kg.
W. rape: Bienvenu, sown at 8 kg.
W. peas: Frijaune, sown at 220 kg.
W. wheat: Avalon, sown at 200 kg.
S. beans: Minden, sown at 220 kg.
S. lupins: Vladimir, sown at 250 kg.
S. peas: Progeta, sown at 230 kg.
Sunflowers: Asmer 077 sown at 80,000 seeds per hectare.

Cultivations, etc.:-

Fallow only: Glyphosate applied: 16 Sept, 1985. K applied, deep-tine cultivated: 18 Sept. Ploughed: 11 Nov. Spring-tine cultivated: 9 May, 1986. Rotary cultivated: 25 June. Deep-tine and rotary cultivated: 13 Aug.

All winter crops: Glyphosate applied: 16 Sept, 1985. K applied, deep-tine cultivated: 18 Sept. Deltamethrin applied: 30 Oct.

W. beans only: Rotary cultivated, seed sown, spring-tine cultivated: 4 Oct, 1985. Weedkillers applied: 11 Oct. Pirimicarb applied: 11 July, 1986. Combine harvested: 11 Sept.

W. oats and w. wheat: Rotary cultivated, seed sown, spring-tine cultivated: 4 Oct, 1985. Weedkiller applied: 11 Oct. N applied: 9 Apr, 1986. Combine harvested: 12 Aug (w. oats), 14 Aug (w. wheat).

W. rape only: Rotary cultivated, seed sown: 18 Sept, 1985. First N applied: 19 Sept. Weedkillers applied: 22 Oct. Second N applied: 11 Mar, 1986. Desiccant with wetting agent applied: 11 Aug. Combine harvested: 13 Aug.

W. peas only: Ploughed: 11 Nov, 1985. Rotary harrowed twice, seed sown: 12 Nov. Pirimicarb applied: 11 July, 1986. Harvested by hand: 28 July.

All spring crops: Glyphosate applied: 16 Sept, 1985. K applied: 18 Sept. Ploughed: 11 Nov. Spring-tine cultivated, seed sown: 8 May, 1986.

S. beans and s. peas only: Pirimicarb applied: 11 July, 1986. Combine harvested: 1 Sept (s. peas), 2 Oct (s. beans).

S. lupins only: Fungicide applied: 10 July, 1986. Harvested by hand and combine threshed: 19 Nov.

Sunflowers only: N applied: 29 May, 1986. Bird repellent applied: 13 Aug. Harvested by hand and combine threshed: 8 Oct. Previous crops: Potatoes 1984, s. barley 1985.

NOTE: Oat and wheat plots were assessed for take-all at the beginning of July.

86/R/CS/320

VARIOUS CROPS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

CROP	
W BEANS	3.76
W OATS	4.83
W RAPE	2.58
W PEAS	0.07
W WHEAT	6.81
S BEANS	4.91
S LUPINS	2.60
S PEAS	3.31
SNFLOWER	1.76
MEAN	3.40

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CROP
-----	-----
SED	0.463

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	16	0.568	16.7

GRAIN MEAN DM% 67.9

PLOT AREA HARVESTED

W BEANS, W PEAS, S PEAS, S BEANS, S LUPINS	0.00115
W OATS, W RAPE, W WHEAT, SNFLOWER	0.00125