

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1986

[Full Table of Content](#)



86/R/PE/1 Effects of Pea Seed-borne Mosaic Virus - Peas

Rothamsted Research

Rothamsted Research (1987) *86/R/PE/1 Effects of Pea Seed-borne Mosaic Virus - Peas* ; Yields Of The Field Experiments 1986, pp 256 - 256 - DOI: <https://doi.org/10.23637/ERADOC-1-36>

86/R/PE/1

PEAS

EFFECTS OF PEA SEED-BORNE MOSAIC VIRUS

Object: To study the transmission, symptoms and effects on yield of pea seed-borne mosaic virus in different varieties of peas with and without insecticide - Fosters West.

Sponsor: A.J. Cockbain.

Design: 3 randomised blocks of 2 whole plots split into 6.

Whole plot dimensions: 20.0 x 5.0.

Treatments: All combinations of:-

Whole plots

1. INSCTCDE	Insecticide:
NONE	None
PP 321	'PP 321' applied at 0.20 l in 220 l on 19 June, 1986, 30 June and 11 July

Sub plots

2. VARIETY	Varieties:
BIRTE	Birte, sown at 230 kg
MARO	Maro, sown at 300 kg
PROGRETA	Progreta, sown at 230 kg
VEDETTE	Vedette, sown at 260 kg
WAVEREX	Waverex, seed infected pea seed-borne mosaic virus (duplicated), sown at 140 kg

- NOTE: (1) The positions of the sub plots of Waverex were allocated to ensure that the other varieties were always adjacent to Waverex.
- (2) Plots were netted against bird damage from the end of July until harvest.
- (3) A strip of s. beans was sown at one end of the experiment and a strip of lupins at the other to assess virus spread to these crops.

Basal applications: Weedkillers: Simazine at 0.17 kg and trietazine at 1.2 kg in 200 l.

Cultivations, etc.:- Heavy spring-tine cultivated: 12 Sept, 1985.
Ploughed: 6 Dec. Spring-tine cultivated, seed sown: 7 May, 1986.
Weedkillers applied: 16 May. Combine harvested: 9 Sept. Previous crops: W. wheat 1984 and 1985.

- NOTES: (1) Aphid numbers were assessed during the growing season. Virus incidence was assessed in the plants at the end of flowering and in seed from some plots after harvest.
- (2) Because of a harvesting error one plot was lost, with treatment combination PP 321, PROGRETA. An estimated value was used in the analysis.